

Pollutants Biodegradation by Fungi

C. Pinedo-Rivilla, J. Aleu and I. G. Collado*

Departamento de Química Orgánica, Facultad de Ciencias, Universidad de Cádiz, s/n, Apdo. 40, 11510 Puerto Real, Cádiz, Spain

Abstract: One of the major problems facing the industrialized world today is the contamination of soils, ground water, sediments, surfacewater and air with hazardous and toxic chemicals. The application of microorganisms which degrade or transform hazardous organic contaminants to less toxic compounds has become increasingly popular in recent years. This review, with approximately 300 references covering the period 2005-2008, describes the use of fungi as a method of bioremediation to clean up environmental pollutants.

1. INTRODUCTION

Pollution of the environment has been one of the largest concerns to science and the general public in the last years. Nowadays, the industrialized world is confronted with the contamination of soils, water sources and air with hazardous and toxic xenobiotics. While regulatory steps have been implemented to reduce or eliminate the production and release to the environment of these chemicals, significant environmental contamination has occurred in the past and will probably continue to occur in the future. The industrialization of agriculture, rapid growth in the chemical industry and the need to generate cheap forms of energy have all caused the continuous release of very organic chemicals into the biosphere. For example, in the United States alone an enormous amount waste is produced annually. In fact, approximately 300 million metric tons of hazardous wastes are produced each year.

Bioremediation is a process by which living organisms degrade or transform hazardous organic contaminants to less toxic compounds [1]. Microorganisms in the indigenous environment have been known to play key roles in the biodegradation of organic compounds. Unlike prokaryotes, eukaryotic fungi have shown diverse metabolic potential resulting in metabolites similar to those produced from mammalian metabolism. These metabolic properties may help us to directly elucidate the metabolic fates of organic compounds occurring in mammalian liver cells instead of using mammalian microsomal fractions or live organisms. Fungal metabolism also provides an easy preparative method for the production of metabolites in large quantity.

The use of fungi as a method of bioremediation provides an option to clean up environmental pollutants. Bioremediation using fungi has drawn little attention in the past two decades since most bioremediation research has focused mainly on the use of bacteria. Nevertheless, recently fungi have received considerable attention for their bioremediation potential which is attributed to the enzymes they produce. In addition, fungi have advantages over bacteria such as fungal

hyphae that can penetrate contaminated soil to reach the pollutants [2]. This review, with approximately 300 references covering the period 2005-2008, will highlight the main applications of fungi to the biodegradation of organic contaminants to less toxic compounds in order to clean up environmental pollutants.

2. BIODEGRADATION OF CHEMICAL POLLUTANTS BY FUNGI

The revolutionized development of resources and technologies has produced more chemicals and compounds which has consequently increased the number of compounds identified as being potential environmental threats to living organisms. Pharmaceuticals and personal care products (PPCPs), surfactants, various industrial additives and numerous chemicals are purported to be pollutants. These pose challenges to the designers of future treatment plants and related methodology for their eradication [3].

These pollutants vary greatly in their form and mechanism of action. Thus, the identification and evaluation of these compounds from the environmental matrixes have provided a unique challenge. The methodologies used for degradation include biological and instrumental methods. The new advances in molecular biology and the isolation of new microorganisms from contaminated environments form the basis of bioremediation emerging as a clean and low-cost methodology for the future.

Aromatic Hydrocarbons

The biodegradation of aromatic hydrocarbons by fungi has traditionally been considered to be of a cometabolic nature. Recently, however, an increasing number of fungi isolated from air biofilters exposed to hydrocarbon-polluted gas streams have been shown to assimilate volatile aromatic hydrocarbons as the sole source of carbon and energy. The biosystematics, ecology, and metabolism of such fungi were reviewed, based in part on the re-evaluation of a collection of published hydrocarbon-degrading isolates obtained from authors around the world [4]. For example, the degradation performance of benzene, toluene, styrene and xylene by fungi was widely studied [5, 6]. Moreover, biodegradation of monomeric styrene by *Phanerochaete chrysosporium* KFRI

*Address correspondence to this author at the Departamento de Química Orgánica, Facultad de Ciencias, Universidad de Cádiz, s/n, Apdo. 40, 11510 Puerto Real, Cádiz, Spain; Tel: 00 34 956 016368; Fax: 00 34 956 016193; E-mail: isidro.gonzalez@uca.es

20742, *Trametes versicolor* KFRI 20251, and *Daldinia concentrica* KFRI 40-1 was carried out by Lee *et al.* [7], giving metabolites including 2-phenyl ethanol, benzoic acid, cyclohexadiene-1,4-dione, butanol and succinic acid.


Oil pollution has become an environment problem which has been paid wide attention in the world. Bioremediation technology applied to oil contaminated soil has become an important field in research for its advantages such as low cost, little environmental effect, simplicity and efficiency and has a bright perspective for the future. The main research fields relating to the bioremediation of oil contaminated soil were put forward employing fungi [8] such as *Trichoderma* sp. [9]. Thus, naphthalan petroleum was studied using fungi cultures to obtain degrader strains as *Penicillium* sp. 3n, *Fusarium* sp. 11a, *Cephalosporium* sp. 45a, and *Mucor* sp. 16 [10, 11]. Also, the fungus *Cladosporium* proved to have good aromatic-degrading ability to biodegrade diesel pollution in aqueous solution [12] and crude oil contaminated soils were used to test the degradation ability of *Pleurotus tuber-regium* where reduction of aromatics was appreciable in all the experiments [13]. Moreover, fuel oil A contaminated sites were investigated for degradation by *Fusarium solani* SZFWT02 showing a great biodegradation activity [14].

Polycyclic aromatic hydrocarbons (PAHs) are toxic pollutants that have accumulated in the environment due to a variety of anthropogenic activities. Bioremediation using various microorganisms is one of the approaches tested for the removal of PAHs from the environment. Fungi belonging to the genera *Aspergillus*, *Penicillium*, *Paecilomyces*, *Coriolus*, *Pycnoporus*, *Pleurotus*, *Fomitopsis*, and *Daedalea*, have been found to be responsible for degrading PHAs in soil and aquatic environments [15]. The degradation potential of white rot fungi belonging to the genera *Phanerochaete*, *Irpex*, *Polyporus*, *Stereum*, *Lentinus*, *Bjerkandera*, *Irpex*, *Pleurotus*, and *Phlebia* to remediate contaminated soils [16-22] is known. The most effective biodegradation of pyrene was obtained with *Coriolus versicolor*, *Trichoderma* sp., *Aspergillus niger*, and *Fusarium* sp. [1, 23]. Other lignolytic fungi, *Allescheriella* sp. strain DABAC 1, *Stachybotrys* sp. strain DABAC 3, and *Phlebia* sp. strain DABAC 9 were selected for remediation of naphthalene, dichloroaniline isomers, *o*-hydroxybiphenyl and 1,1'-binaphthalene [24]. For anthracene (AC) degradation, *Tetrahymena pyriformis* accumulated high amounts of AC without any transformation. In contrast, the fungi *Absidia cylindrospora*, *A. fusca*, *Cunninghamella elegans*, *Aspergillus terreus*, *Cladosporium herbarum*, *Penicillium chrysogenum*, *Rhodotorula glutinis*, and *Saccharomyces cerevisiae*, were able to transform AC to 1,4-dihydroxyanthraquinone as a product of biotransformation [25]. Biodegradation of phenanthrene by *Trichoderma* sp. S019 afforded 1-hydroxy-2-naphthoic acid, salicylaldehyde, salicylic acid, and catechol as intermediates in the bioremediation process [26]. In addition, *Cunninghamella elegans* IM 1785/21Gp gave the metabolites *trans*-1,2,3,4- and 9,10-dihydrodiols, phenols, diphenols (diols), and glycoside conjugates of 1-,2-,3-,4-, and 9-phenanthrols [27]. Other experiments based on co-cultures were carried out with two fungi, *Aspergillus terreus* and *Penicillium* sp., and the bacterial strain *Rhodococcus* sp. IC10 [28]. Moreover, using

Fusarium solani, a high degradation of phenanthrene was obtained in free cultures and immobilized [29, 30, 31].

In the meanwhile, Li *et al.* [32] reported the biodegradation of 1,2,3,4-tetrahydronaphthalene (THN) by the marine fungus *Hypoxylon oceanicum* (326#) giving one major product, 3,4-dihydro-4-hydroxy-1(2*H*)-naphthalenone, and three minor products: 3,4-dihydro-1(2*H*)-naphthalenone, 1,2,3,4-tetrahydro-1-naphthalenol, and 1,2,3,4-tetrahydro-1,2-naphthalenediol.

Other PAHs were tested for fungi biodegradation such as benzo(α)anthracene, benzo(α)fluoranthene, benzo(α)pyrene (**1**), and chrysene, which were biodegraded by *Fusarium flocciferum*, *Trichoderma* species, *Tremetes versicolor*, and *Pleurotus ostreatus* [33]. Benzo(α)pyrene (BaP) (**1**) is a 5-ring polycyclic aromatic hydrocarbon and a large number of fungi were tested for its degradation such as *Trichoderma* sp., *Aspergillus niger*, *Mucor* sp., and *Fusarium* sp. [19, 34, 35, 36]. *Fusarium* sp. E033 was able to biodegrade 65-70% of the initial benzo(α)pyrene (**1**) provided giving two transformation products, dihydroxy dihydro-benzo(α)pyrene and benzo(α)pyrene-quinone [37]. Meanwhile, *Penicillium chrysogenum* SF04 had the highest degradation of BaP (**1**) (up to 71.31 %) [38]. Furthermore, benzo(α)anthracene was degraded by *Irpex lacteus* affording 2-hydroxymethyl benzoic acid or monomethyl- and dimethyl-esters of phthalic acid and 1-tetralone as final products [39]. However, a high degree of benzo(α)pyrene (**1**) degradation is undesirable for the bioremediation of BaP-contaminated soils because some of its accumulated metabolites still have severe health risks for humans such as benzo(α)pyrene-1,6-quinone (BP1,6-quinone) (**2**) and 3-hydroxybenzo(α)pyrene (3-OHBP) (**3**) (Scheme 1) [40].



Scheme 1. Degradation of BaP (**1**) by fungi.

Other groups of important aromatic contaminants are phenols and derivatives. The major sources of phenol contamination are the chemical and petrochemical industries, agriculture (pesticides, containing hydrocarbons), wood processing as part of papermaking technologies, textile industry, etc. The ubiquitous nature of phenols, their toxicity even in trace amounts and the stricter environmental regulations make it necessary to develop processes for the removal of phenols from wastewaters. Biodegradation allows for the utilization of aromatic hydrocarbons by the biological agent and for their re-entrance into the carbon cycle [41].


Along these lines, white-rot fungi have been shown to exhibit unique biodegradation capabilities for phenols [42, 43, 44]. For instance, the fungus *Panus tigrinus* CBS 577.79 was investigated for its ability to reduce the polluting load of olive-mill wastewater (OMW) with a significant presence of phenolic components [45].

On the other hand, strains from *Fusarium* were able to reduce aromatic components by 65% in olive-mill dry residue (DOR) [46]. Moreover, *Fusarium* sp. HJ01 was able to grow using phenol as the only carbon resource giving the intermediate catechol as a biotransformation product [47, 48]. The fungus *Trametes versicolor* was capable of decolouring and degrading phenol compounds from paper mill effluent [49, 50, 51].

As the symbiotic fungi and plant, mycorrhiza was able to degrade organic pollutants [52]. Some conifer ectomycorrhizae can degrade and detoxify water-solution phenolic compounds produced by the conifer *Kalmia angustifolia*. Thus, *Paxillus involutus*, *Laccaria laccata*, and *L. bicolor* were employed to degrade water leachates of *Kalmia* leaf and litter. Pure ferulic, *o*-coumaric, and *o*-hydroxyphenylacetic acids were degraded by 100, 98, and 79.5%, respectively, in the presence of *P. involutus* 211804 [53].

The biodegradation of polychlorophenols (PCBs) [54], an important group of phenols which have been used as fungicides, herbicides, insecticides, and in the synthesis of other pesticides, has been widely studied [55-58]. The white-rot fungus *Phlebia brevispora* was shown to be able to degrade PCBs obtaining *m*-methoxylated, *p*-dechlorinated and *p*-methoxylated metabolites [59]. 4,4'-Dichlorobiphenyl (4,4'-DCB) and its metabolites were added to cultures from *Phanerochaete* sp. and the metabolic pathway was elucidated by the identification of metabolites namely 2-hydroxy-4,4'-DCB and 3-methoxy-4,4'-DCB, 4-chlorobenzoic acid, 4-chlorobenzaldehyde, 4-chlorobenzyl alcohol, and 4-hydroxy-3,4'-DCB [60, 61, 62]. Fungi *Phlebia* sp., *Phanerochaete chrysosporium*, and *Mortierella* sp. were also selected to degrade different chlorobenzoic acids (CBA) [63] giving some aromatic metabolites from a hydroxylation pathway and a dechlorination pathway [64]. Furthermore, *Bjerkandera adusta*, *Anthracoephyllum discolor*, immobilized and non-immobilized *Phanerochaete chrysosporium* [65], *Trametes versicolor* isolate HR131, and *Trametes* sp. isolate HR577 were studied in the degradation of pentachlorophenol [51, 66, 67, 68]. Fungi *Boletus edulis*, *Suillus luteus*, *Cortinarius russus*, *Suillus grevillei*, *Gomphidius viscidus*, *Laccaria bicolor*, *Leccinum scabrum*, *Xerocomus chrysenteron*, *Heboloma crustuliniforme*, and *H. longicaudum* were grown in media with different substrate concentrations of pentachlorophenol (PCP) to determinate their effect on fungal growth. No impact on the growth of the mycelia was observed at low ambient PCP levels. In addition, the high tolerance capability for pentachlorophenol may be related to their oxidoreductase activities and acidification effect [69].

Whereas, *Aspergillus awamori* NRRL 3112 degraded phenol, catechol, 2,4-dichlorophenol and 2,6-dimethoxyphenol as well as *Peniophora cinerea*, *Psilocybe castanella*, two strains of *Trametes villosa*, *Agrocybe perfecta*, *Trichaptum bisogenum* and *Lentinus villosus* were able to colonize soil containing up to 4600 mg pentachlorophenol/kg




Scheme 2. Fungal biotransformation products of dehydroabiestic acid (6).

soil. All fungi produced chloride ions during degradation, indicating dehalogenation of the molecule [70]. Moreover, Taseli *et al.* [71] studied the potential of the fungus *Penicillium camemberti*, which degraded pentachlorophenol (PCP), 2-chlorophenol and trichloroacetic acid.

Other phenolic compounds with special activities are nonylphenols (4) and bisphenol A (5) (BPA), known as endocrine-disrupting compounds. Technical nonylphenol (t-NP) mixtures (4) were assessed using the mitosporic fungal strain UHH 1-6-18-4 and a strain of the aquatic hyphomycete *Clavariopsis aquatica*. All t-NP isomers were degraded to individual extents [72]. On the other hand, Soares *et al.* [73] showed that the fungi *Phanerochaete chrysosporium*, *Pleurotus ostreatus*, *Trametes versicolor* and *Bjerkandera* sp. BOL13 degraded nonylphenol (4) at an initial concentration of 100 mg/L. In addition, bisphenol A (5) was biodegraded with several white rot fungi (*Irpex lacteus*, *T. versicolor*, *Ganoderma lucidum*, *Polyporellus brumalis*, *Pleurotus eryngii*, *Schizophyllum commune*) isolated in Korea and two transformants of *T. versicolor* (strains MrP 1 and MrP 13) [74]. *Stereum hirsutum* and *Heterobasidion insulare* showed high resistance to BPA (5) [75].

Further results showed the potential of the fungi *Basidi-radulum molare* and *Schizopora paradoxa* to degrade phenolic compounds such as 4-*tert*-octylphenol [76]. Moreover, van Beek *et al.* [77] reported the degradation of dehydroabiestic acid (DHA) (6) from Scots pine wood by *Trametes versicolor* and *Phlebiopsis gigantea* in liquid stationary cultures, isolating some biodegradation products from *P. gigantea* cultures: 1 β -hydroxy-DHA (7), 1 β ,7 α -dihydroxy-DHA (8), 1 β ,16-dihydroxy-DHA (9), and tentatively 1 β -hydroxy-7-oxo-DHA (10) and *T. versicolor* cultures, 1 β ,16-




Scheme 3. Biodegradation of the polychlorinated dioxins **17-20**.


dihydroxy-DHA (**9**), 7 β ,16-dihydroxy-DHA (**11**), 1 β ,7 β ,16-trihydroxy-DHA (**12**), 1 β ,16-dihydroxy-7-oxo-DHA (**13**), 1 β ,15-dihydroxy-DHA (**14**), and 1 β ,7 α ,16-trihydroxy-DHA (**15**) (Scheme 2). Also, *Candida tropicalis* was tested on the reduction of free gossypol (**16**), a polyphenol derived from the cotton plant. This biodegradation was evaluated through optimization of the parameters in order to operate under optimal conditions [78].

Indeed, polychlorinated dibenzo-*p*-dioxins (PCDDs) and polychlorinated dibenzofurans (PCDFs), commonly known as dioxins (PCDD/Fs) [79], are toxic environmental pollutants formed from various sources. Removal of dioxins by biological degradation is considered a feasible method as an alternative to other expensive physic-chemistry approaches [80]. Different dibenzo-*p*-dioxins and dibenzofurans (PCDD/Fs) were studied for degradation by fungi such as *Phlebia radiata* I-5-6 [81, 82, 83], *P. acerina*, *P. lindtneri* and *P. brevispora* which can hydroxylate and methoxylate PCDDs [84], and *Phanerochaete chrysosporium* DSM 6909, *P. chrysosporium* DSM 1556, *Irpex* sp. W3, *Trametes* sp. CH2, *Fusarium* sp. VSO7 [85], and *Pleurotus pulmonarius* [86]. Biodegradation of 2,8-dichlorodibenzo-*p*-dioxin (2,8-DCDD) (**17**), 2,3,7,8-tetrachlorodibenzo-*p*-dioxin (2,3,7,8-TCDD) (**18**), 2,7-dichlorodibenzo-*p*-dioxin (2,7-DCDD) (**19**), and 2,4,8-trichlorodibenzofuran (2,4,8-TCDF) (**20**) was conducted with two fungi (PL1 and 267) already screened from nature [87]. Furthermore, 2,8-DCDD (**17**) and 2,3,7,8-TCDD (**18**) were also degraded by those fungi, giving compounds considered to be intermediates, namely 4-chlorocatechol (**21**) and 4,5-dichlorocatechol (**22**) respectively [88, 89]. Chlorocatechol (**21**), 3,5-dichlorosalicylic acid (**23**) and 5-chlorosalicylic acid (**24**) were isolated from 2,7-DCDD (**19**) and 2,4,8-TCDF (**20**), respectively (Scheme 3) [90].

Lastly, other aromatic pollutants have been tested for bioremediation. Quinoline (**25**) was biodegraded by *Pleurotus ostreatus* BP resulting in total mineralization and some fermentation products [91].

As the use of hydrocarbons by the microorganisms is associated with biosurfactant production, biodegradation by *Aspergillus niger* and *Penicillium nigricans* was also investigated [92]. Another *A. niger* strain namely PSH is capable of degrading tannins giving gallic acid (**26**) and ellagic acid (**27**) [93]. Martin *et al.* [94] reported the degradation of galaxolide (HHCB) (**28**) and tonalide (AHTN) (**29**), two micro-pollutants from aquatic environments, by *Myrioconium* sp. strain UHH 1-13-18-4 and *Clavariopsis aquatica*. The products obtained were the result of hydroxylations at different positions. *Polyporus brumalis* was applied to degrade dibutyl phthalate (DBP), the main product being phthalic acid anhydride as well as trace amounts of α -hydroxyphenylacetic acid, benzyl alcohol, and α -hydroxyphenylacetic acid [95].





Additionally, degradation of refractory organic matter (OM) by the basidiomycete fungus *Schizophyllum commune* and white rot fungi was reported. Main products of the biodegradation were organic heavy metal complexes which can enter the environment [96, 97].

Moreover, indole degradation was studied by *Sporotrichum thermophile* and *Pleurotus ostreatus* with more than a 99% consumption rate of indole [98, 99]. Biodegradation of *p*-cresol by *Gliomastix indicus* was studied [100].

Aliphatic Hydrocarbons

Fungi can also degrade *n*-alkanes such as tridecane, tetradecane, pentadecane, hexadecane, heptadecane, octadecane (C13-C18) and crude Omani oil. Biodegradation by the fungi *Aspergillus niger*, *A. ochraceus*, *Trichoderma asperellum* strain TUB F-1067 (SA4), *T. asperellum* strain Tr48 (SA5), *T. asperellum* strain TUB F-756 (SA6), *Penicillium* species (P1), and *Aspergillus* species (P9) was studied [101]. Among these fungi, the P1 strain exhibited greater potential in degrading the aliphatic hydrocarbon compounds of used motor oil [2].

Some of the most important aliphatic hydrocarbon pollutants are *n*-eicosane, which was degraded by *Trichoderma* sp. S019 affording nonadecanoic acid, *n*-octadecane, hexadecanoic acid, oleic acid and stearic acid as reaction products [102]. Imidazolium compounds (ICs) and quaternary ammonium compounds (QACs) were degraded by two strains of *Gliocladium roseum*, *Penicillium brevi-compactum*, *P. funiculosum*, *Phialophora fastigiata*, *Verticillium lecanii* [103]. Carbon tetrachloride (CT) [104], trichloroethylene (TCE) and perchloroethylene (PCE), one of the most important groundwater pollutants, were tested for degradation by fungi. The aerobic degradation of PCE was reported for the first time by *Trametes versicolor*, giving 2,2,2-trichloroethanol and CO₂ as main byproducts from TCE degradation, and trichloroacetic acid (TCA) from PCE [105]. Moreover, *Ganoderma lucidum* and *Irpex lacteus* were able to degrade substantial levels of perchloroethylene (PCE) and trichloroethylene (TCE) in pure culture [106].

Another white-rot fungi, *Bjerkandera adusta*, was able to degrade hexachlorocyclohexane (HCH) isomers giving 1-(3-chloro-4-methoxyphenyl)ethanone and (2,4-dichloro-3-methoxy)-1-benzenecarbonyl chloride demonstrating the capability of *B. adusta* to produce these types of organochlorine compounds [107].

Compounds with different activities have been investigated for biotransformation including the sesquiterpene botrydiediol (**30**) by *Botrytis cinerea* affording three new sesquiterpenoids [108]. Wang *et al.* [109] studied the biodegradation by several fungi of digitoxin (**31**), a cardiac glycoside that is presumed to be effective in the treatment of heart failure. *Curvularia lunata* AS3.3589 and *Absidia coerulea* CICC40302 gave some biotransformation products.

Cyanide Compounds

Organic and inorganic cyanide compounds are widely distributed on the planet and they are among the most common corrosive pollutants. Most people associate the word cyanide with an extremely dangerous and fast-acting poison. However, there are several cyanide species, of varying toxicity, depending on the source of cyanide contamination. Free cyanide is the most toxic form and is easily and rapidly absorbed through inhalation, ingestion or skin contact. Thiocyanates are much less toxic than free cyanide and iron-complexed cyanides are only mildly toxic [110].


The degradation of simple cyanides has also been demonstrated in fungi [111, 112]. A fungal mutant of *Trichoderma koningii*, TkA8, constructed by restriction enzyme-mediated integration, was shown to have a high cyanide degradation ability [113]. As shown by Hossain *et al.* [114], *Trametes versicolor* ATCC 200801, *Phanerochaete chrysosporium* ME 496 and *Pleurotus sajorcaju* tolerated up to 500-ppm initial concentration of cyanide.

3. PESTICIDES



The agricultural industry's dependency on chemicals to sustain productivity in marginal landscapes has led to a global-scale contamination of the environment with toxic pesticides and nutrient fertilizers which are changing the course of biogeochemical cycles. They include fungicides, insecticides, and herbicides and are one of the causes of water pollution, and some pesticides are persistent organic pollutants contributing to soil contamination. Among the techniques employed to remove these contaminants, biodegradation is very effective, less contaminating and cheaper than others.

Fungicides

Biphenyl (**32**) and the monohydroxylated derivatives 2-hydroxy- and 4-hydroxybiphenyl are known to be fungistatic substances. These compounds are widely used for the con-



Scheme 4. Biodegradation of biphenyl (**32**) by *Talaromyces elicus*.



servation of citrus fruits, even though biphenyl (**32**) is known for its toxic effects on humans. The filamentous fungus *Talaromyces helicus* oxidized biphenyl (**32**) to the hydroxylated derivatives 4,4'-dihydroxybiphenyl (**33**), 3,4-dihydroxybiphenyl (**34**), 2-hydroxybiphenyl (**35**), 2,5-dihydroxybiphenyl (**36**), and the ring cleavage product 4-phenyl-2-pyrone-6-carboxylic acid (**37**) (Scheme 4) [115].

Tribromophenol (TBP) is used in wood preservation. *Trametes versicolor* and *Agaricus augustus* proved effective in decreasing TBP concentrations and *A. augustus* was also capable of biotransforming TBP to tribromoanisole (TBA) [116]. Other fungi employed for the degradation of this fungicide were *Laetoporeus sulfureus*, *Gloeophyllum trabeum*, and *Ganoderma australe* in liquid culture, and were able to degrade TBP, degradation by *G. australe* being the most efficient (71% to 77%) [117].


In addition, widdrol (**38**) has shown activity against the necrotrophic plant pathogen *Botrytis cinerea*. The biotransformation of **38** by *B. cinerea* and *Colletotrichum gloeosporioides* afforded four and one biotransformation products (**39-43**), respectively. Biotransformation with *C. gloeosporioides* yielding for the most part oxidation products at C-10: 10-oxowiddrol (**39**), 10 β -hydroxywiddrol (**40**), 10 α -hydroxywiddrol (**41**), and 14 α -hydroxywiddrol (**42**). The biotransformation products were then tested against *B. cinerea* and found to be inactive [118].

Another group of pesticides with fungicide activity is tributyltin (**44**) compounds, a group of compounds containing the $(\text{C}_4\text{H}_9)_3\text{Sn}$ moiety. The filamentous fungus *Cunninghamella elegans* was able to degrade tributyltin chloride (TBT) giving less toxic compounds, dibutyltin and monobutyltin [119, 120].


Insecticides

Some insecticides have been degraded by several fungi. For instance, endosulfan (**45**), widely employed as pesticide (insecticide and acaricide), was degraded by *Chaetosartorya stromatoides*, *Aspergillus terricola*, and *A. terreus* showing degradation rates of up to 75% [121]. In other studies using *A. niger*, various intermediates of endosulfan (**45**) metabolism including endosulfan diol and endosulfan sulfate were isolated [122].

Romero *et al.* [123] reported the biodegradation of toxaphene (**46**) in waste substrates by the fungus *Bjerkandera* sp. strain BOL13. One of the most important insecticides that is extensively used and toxic is lindane (**47**) [124], which was tested for biodegradation by nonwhite rot fungi [125] and white-rot fungi such as *Phanerochaete chrysosporium*, *Trametes hirsutus*, *Bjerkandera adusta*, and *Pleurotus* sp. [126, 127]. Also imidacloprid (**48**), a class of neuro-active insecticide modeled after nicotine, was degraded by *Calocybe indica* [128].



Scheme 5. Biodegradation of carbofuran (**49**) by *Mucor ramannianus*.



Scheme 6. Biodegradation of *N,N*-diethyl-*m*-toluamide (**53**).

Another example of insecticide biodegraded by several fungi is carbofuran (**49**), which was added to cultures of *Gliocladium* [129] and *Mucor ramannianus* affording 2-hydroxy-3-(butan-2-ol)phenol (**50**) and 3-hydroxycarbofuran-7-phenol (**51**) as transformation products (Scheme 5) [130].

Brown-rot fungi were also investigated for their ability to degrade 1,1,1-trichloro-2,2-bis (4-chlorophenyl)ethane (DDT), as well as white-rot fungi [131] and ectomycorrhizal fungi [132]. For instance, *Gloeophyllum* genus, *Daedalea* genus, and *Fomitopsis* genus showed a high ability to degrade DDT affording 1,1-dichloro-2,2-bis (4-chlorophenyl) ethane (DDD), 1,1-dichloro-2,2-bis(4-chlorophenyl)ethylene (DDE), and 4,4-dichlorobenzophenone (DBP) as metabolic products [133].

β -Cyfluthrin (**52**), a synthetic pyrethroid insecticide, was biotransformed by *Aspergillus nidulans* and *Sepedonium maheswarium* to 4-fluoro-3-phenoxybenzaldehyde and 3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropane carboxylic acid [134].

Seo *et al.* [135] reported the biotransformation of *N,N*-diethyl-*m*-toluamide (DEET) (**53**), a topical insect repellent, by *Cunninghamella elegans* ATCC 9245, *Mucor ramannianus* R-56, *Aspergillus niger* VKMF-1119, and *Phanerochaete chrysosporium* BKMf-1767 to *N,N*-diethyl-*m*-


toluamide-*N*-oxide (**54**), *N*-ethyl-*m*-toluamide-*N*-oxide (**55**), and *N*-ethyl-*m*-toluamide (**56**) (Scheme 6).

The biodegradation by the strain *Aspergillus nomius* L3 of the insecticide dimethoate (**57**) as a cosubstrate was reported by Ai *et al.* [136]. In addition, methoxychlor (**58**) was converted to 2,2-dichloro-1,1-bis(4-methoxyphenyl)ethane, 2,2-dichloro-1,1-bis(methoxyphenyl)ethylene, 2-chloro-1,1-bis(4-methoxyphenyl)ethane, 2-chloro-1,1-bis(4-methoxyphenyl)ethylene, and 1,1-bis(4-methoxyphenyl)ethylene by *Stereum hirsutum* [137].

Lastly, Osman *et al.* [138] studied the biotransformation of dicofol pesticide (DCF) (**59**) by *Trichoderma viride* and *T. harzianum* with no intermediate or final degradation metabolites isolated. Fenitrothion breakdown product 3-methyl-4-nitrophenol (MNP), a newly characterized estrogenic chemist, was biotransformed by *Aspergillus niger* VKM F-1119 to 2-methyl-1,4-benzenediol, 4-amino-3-methylphenol, and two singly hydroxylated products [139].

Herbicides

Herbicides are widely used in agriculture to kill undesired plants. However, because of the large number of herbicides in use, there is significant concern regarding health effects which make their elimination from soil and water, an important focus of study.



Scheme 7. Biodegradation of isoproturon (**63**) by fungi.


For example, ectomycorrhizal fungi from rice were evaluated for their applicability in studies of herbicide degradation. One of those strains was studied for its ability to degrade propanil (**60**) and its metabolite 3,4-dichloroaniline (3,4-DCA) [140]. Other fungi capable of degrading herbicides such as sulfentrazone (**61**) were *Chrysosporium* sp., *Eupenicillium* sp., and *Paecilomyces* sp. [141]. Romeh *et al.* [142] investigated the degradation of fluometuron (**62**) herbicide by *Trichoderma viride*, *Metarhizium anisopliae*, and *Beauveria bassiana* showing biodegradation rates up to 85%.

Other herbicides were also employed to study their biodegradation by fungi, such as isoproturon (**63**), which was converted by *Phoma eupyrena*, *Mucor hiemalis*, and *Mortierella* sp. to hydroxylated metabolites *N*-(4-(2-hydroxy-1-methylethyl)phenyl)-*N'*,*N'*-dimethylurea (**64**) and *N*-demethylated metabolite *N*-(4-isopropylphenyl)-*N'*-methylurea (MDIPU) (**65**) (Scheme 7) [143]. Chlornitrofen (CNP) (**66**)

was studied for biodegradation by *Phlebia brevispora* TMIC33929. In the degradation experiment using CNP (**66**) standard compounds, CNP (**66**) was transformed into several metabolites including monomethoxylated compounds and 2,4,6-trichlorophenol [144]. Another example of herbicide is glyphosate (**67**), a systemic non-selective herbicide, which was degraded by *Fusarium* sp. [145]; Zhu *et al.* [146] investigated the biodegradation by fungi of acetochlor (**68**) in soil. The effects of metsulfuron-methyl (**69**), a sulfonylurea herbicide, on soil microorganisms were evaluated in various experiments showing that fungi such as *Penicillium* sp. were highly tolerant [147, 148].

Metals

Soil contamination by toxic metals has become a serious problem, because of their long-term persistence and their diffusion into underground water. Heavy metal and non-



degradable chemical contamination of soil and water is a major environmental threat. In recent years, worldwide researchers are investigating new sustainable methods to mitigate such environmental contamination, such as biodegradation by fungi.

The common filamentous fungi can absorb heavy metals (Zn, Cd, Pb, Fe, Ni, Ag, Th, Ra & U) from aqueous solutions. The availability of a variety of fungi with different characteristics and metal binding potential makes it an economical and sustainable option for the removal and recovery of heavy metals [149].

Contaminated soil containing sulfide ore ashes and aromatic hydrocarbons could be treated with a metal-resistant strain BAS-10 of *Klebsiella oxytoca* and other fungi added to the soil, *Allescheriella* sp. DABAC 1, *Stachybotrys* sp. DABAC 3, *Phlebia* sp. DABAC 9, *Pleurotus pulmonarius* CBS 664.97, and *Botryosphaeria rhodina* DABAC P82. *B. rhodina* was the most effective fungus leading to the depletion of the most abundant contaminants [150].

Baldi *et al.* [151] developed a novel process combining sequential treatments of contaminated soil from the ACNA site (Cengio, Savona, Italy). The soil was leached to remove metals in the following order: Pb (74.2%) > Cu (72.6%) > Zn (40.2%) > Ni (55.7%) > Cd (41.5%) > Cr > (21.7%) Co > (19%) Fe (8.2%). The leachate was then incubated with the metal-resistant *Klebsiella oxytoca* strain BAS-10 and *Allescheriella* sp. DABAC 1 leading to a complete degradation of several organic contaminants.

Several metal ions were bioremediated by fungi. For instance, the treatment of the simulated lead-contaminated solid waste by composting with white-rot fungus was studied in the laboratory [152]. The white-rot basidiomycete *Phanerochaete chrysosporium* was very effective in the bioremediation of Pb-contaminated soil [153]. The Pb(II) biosorption potential of *Aspergillus parasiticus* [154] and the macrofungus *Ganoderma carnosum* [155] were also studied in a batch system and biosorption conditions were optimized showing that biosorption potential depended on physicochemical parameters. Tunali *et al.* [156] reported the biosorption of Pb(II) onto *Cephalosporium aphidicola* and the nature of the possible cell and metal ion interactions was examined by the FTIR technique. Moreover, the potential of *Botrytis cinerea* as a biosorbent for metal ions such as Zn(II) and Pb(II) was studied. Competitive biosorption experiments were performed with Zn(II) in the presence of Cu, Cd, and Ni ions simultaneously [157], demonstrating that other competing metal ions (Cu(II), Cd(II), and Ni(II) co-cations) reduced the biosorption capacity on Pb(II) and Zn(II) [158]. *Aspergillus flavus* and *Neurospora crassa* fungal biomass were also able to absorb Pb(II) and Cu(II) under optimum conditions [159, 160]. Cr(VI) biosorption with *Trichoderma viride*, *Aspergillus niger*, *A. sydoni*, and *Penicillium janthinellum* biomass was studied by Bishnoi *et al.* [161] showing that biosorption of Cr(VI) was pH dependent and the maximum adsorption was at pH 2.0. Cr(VI) removal was 91.03% using *A. niger* and 87.95% and 86.61% with *A. sydoni* and *P. janthinellum* [162]. Lastly, the brown-rot fungus *Lentinus edodes* was used as an efficient biosorbent for the removal of Cd from water and three kinds of adsorption models were applied to simulate the biosorption data [163].

Mercury is one of the most harmful ion metals and its biotransformation by fungi was investigated in *Hymenoscyphus ericae*, *Neocosmospora vasinfecta*, and *Verticillium terrestre* following the exposure of these fungi to environmentally relevant doses of Hg (II) (HgCl₂) in aerated pH-controlled cultures [164].

Industrial Dyes







Dyes, originally obtained exclusively from natural sources, are today also produced synthetically on a large scale and represent one of the very mature and traditional sectors of the chemical industry.

Different dyes and pigments are extensively used in the textile, paper, plastic, cosmetics, pharmaceutical and food industries. Most of the earlier dye decolourization studies were based mainly on white-rot fungi such as *Phanerochaete chrysosporium*, *Trametes versicolor*, *Phellinus gilvus*, *Pleurotus sajor-caju*, *Pycnoporus sanguineus*, *Dichomitus squalens*, *Irpex flavus*, *Daedalea flavida*, *Polyporus sanguineus*, *Funalia trogii* ATCC200800, *Ischnoderma resinsum* [165], *Dichomitus squalens* [166], and *Ganoderma* sp [167-170].

Textile industries consume large amounts of water and their effluents contain a wide range of contaminants. These contaminants are dyes with strong colour, inorganic salts, as well as high pH. Textile wastewater containing significant concentrations of dyes cause substantial treatment problems. Most of the dye molecules have a polyaromatic structure with a high molecular weight and contain atoms of nitrogen, sulfur and metals making it very difficult to break them down [171]. Biological methods are emerging as an effective alternative for chemical approaches. For instance, fungi have shown a strong resistance to dye toxicity and it would therefore be a good idea to study fungal strains to identify the potential fungal candidates for dye removal and biodegradation [172-177].

White-rot fungi have been widely employed to biodegrade textile dyes [178-181]. Thus, Sukumar *et al.* [182] investigated the decolourizing ability of *Phanerochaete* sp. and *Trametes* sp., recording colour reduction of 82.01% and 76.07% respectively [183]. The potential of *Trametes villosa* and *Pycnoporus sanguineus* to decolourize reactive textile dyes was evaluated, *T. villosa* being the best degrader [184]. The characteristics of biodegradation of the classical triphenylmethane dyes such as Crystal Violet (**70**), Malachite Green (**71**), and Bromophenol Blue (**72**) were reported by white rot fungi in rice straw media [185, 186, 187]. Yan *et al.* [188] studied the decolourization by biosorption using dead white rot fungus *Pleurotus ostreatus* BP, whose decolourization rate for Remazol Brilliant Blue R (RBBR) (**73**) reached 82.35%.

Synthetic dyes are released in wastewater from textile manufacturing plants, and many of these dyes are genotoxic. For instance, *Irpex lacteus* was used for mutagenicity assays showing that all dyes except Congo Red (CR) (**74**) were mutagenic, indicating that the combined biodegradation process may be useful for reducing the mutagenicity associated with wastewater from textile industries [189]. The biodegradation of Methyl Orange (**75**), Yellow RR Gran, Congo Red (**74**), Bismarck Brown (**76**), Brilliant Red K-2BP (**77**), and the azo dye Remazol Red RR Gran in cultures of the









78: R₁=R₂=R₅=H; R₃=R₄=SO₃Na; R₆=OH
80: R₁=R₃=R₄=R₆=H; R₂=SO₃Na; R₅=OH
82: R₁=R₃=R₄=R₅=H; R₂=SO₃Na; R₆=OH
83: R₁=Me; R₂=SO₃Na; R₃=R₄=R₅=H; R₆=OH

white rot fungus *Phanerochaete chrysosporium* was demonstrated by decolourization studies [190, 191, 192, 193]. Moreover, the ability to decolourize eight chemically different synthetic dyes (Orange G (**78**), Amaranth (**79**), Orange I (**80**), Remazol Brilliant Blue R (RBBR) (**73**), Cupphthalocyanin, Poly R-478 (**81**), Malachite Green (**71**) and Crystal Violet (**70**)) by the white rot fungus *Dichomitus*

squalens was evaluated showing high decolourization capacity for all dyes tested, but not to the same extent [194].

Asgher et al. [195, 196] investigated the indigenous white rot fungi *Pleurotus ostreatus* IBL-02, *Phanerochaete chrysosporium* IBL-03, *Coriolus versicolor* IBL-04, *Ganoderma lucidum* IBL-05, and *Schizophyllum commune* IBL-06 for decolourization of several textile dyes. The results



showed that *P. chrysosporium* could decolourize all the dyes tested, and *C. versicolor* IBL-04 degraded all the dyes, except Drimarene Orange K-GL. Moreover, *P. ostreatus* also showed good decolorization efficiencies on all dyes, except Remazol Brilliant Yellow. However, the rest of the strains showed poor decolorization potential. In addition, *S. commune* IBL-06 and *G. lucidum* IBL-05 were able to degrade Solar Golden Yellow R [197], Solar Orange RSN [198], and Cibacron Red FN-2BL [199].


Moreover, Machado *et al.* [200, 201] studied the potential of basidiomycetous fungi isolated from tropical ecosystems to remove Remazol Brilliant Blue R (RBBR) (**73**) dye. *Trogia buccinalis* showed the highest RBBR (**73**) decolorization.

Removal of water-solution sulfonated azo dyes from textile industry effluents is a major issue in wastewater treatment [199]. Biodegradation of sulfonated azo dyes was studied using white-rot fungi affording 4-hydroxy-benzenesulfonic acid, 3-methyl-4-hydroxy-benzenesulfonic acid, benzenesulfonic acid, 1,2-naphthoquinone-6-sulfonic acid, and 3-methyl-benzenesulfonic acid, as major biotransformation products for Orange I (**80**), Acid Orange 7 (**82**), Acid Orange 8 (**83**), Great Acid Red, and 4-[(4-hydroxyphenyl)azo]-benzenesulfonic acid (**84**) [202, 203, 204]. The degradation of Mordant Violet 5 (**85**) by *Pleurotus ostreatus*

gave benzenesulfonic acid, 4-hydroxybenzenesulfonate, and 1,2-naphthoquinone [205, 206]; Disperse Orange 3 (**86**), Disperse Orange 1 (**87**), and Disperse Red 1 (**88**) were degraded to nitrobenzene, 4-nitrophenol and 4-nitroaniline [207, 208]. Furthermore, 1-methoxy-4-nitrobenzene, 1,2-dimethoxy-4-nitrobenzene, and 2-methoxy-4-nitrophenol were found to be produced from 4-nitrophenol [218].

Decolourizing by non white-rot fungal species such as *Aspergillus flavus*, *A. niger* [209, 210], *Helminthosporium* sp, *Mucor* sp, *Penicillium* sp., *Trichoderma viride*, *Myrothecium* sp. IMER1 [211], and *Fusarium* sp. isolated from textile effluent, was investigated for Remazol Yellow, Remazol Orange [212], Remazol Brilliant Blue R (RBBR) (**73**), Neutral Brilliant Blue GL, and Acid Blue B (**89**), obtaining decolorization rates ~100% [213, 214, 215]. Also, *Trichophyton rubrum* LSK-27 was able to decolourize 83% of Remazol Tiefschwarz, 86% of Remazol Blue RR (**90**) and 80% of Supranol Turquoise GGL in liquid cultures [216].

The mushroom fungi *Lentinus conatus*, *Ischnoderma resinosum*, and *Ganoderma lucidum* KMK2 were studied for this purpose [217]. The results showed that *I. resinosum* was able to decolourize all textile dyes tested (Reactive Black 5 (**91**), Reactive Blue 19 (**73**), Reactive Red 22 (**92**), and Reactive Yellow 15 (**93**)) [218] and *G. lucidum* decolorized anthraquinone dye Remazol Brilliant Blue R (RBBR) (**73**) and



91

diazo dye Remazol Black-5 (RB-5) (**91**) with biodegradation levels up to 90% [219]. Furthermore, *Coprinellus xanthothrix*, a new fungal strain isolated from a polyphenol polluted soil in Greece, was tested for its ability to degrade a polyaromatic dye Poly R-478 (**81**). The fungus showed biosorption and biotransformation as removal mechanisms [220].

Dyehouse effluent treatment has become inevitable because of the presence of dyes which originate from harmful chemicals. *Galactomyces geotrichum* and *Trametes* sp. isolated from contaminated soil were employed to degrade these dyes, finding a new strain with the high rate of 87.21% color reduction [221, 222]. In addition, *Trametes versicolor* decolourized the mono-azo-substituted naphthalenic dye Amaranth (**79**) [223, 224].

Lignin and Cellulose Degradation

Cellulose is the main polymeric component of the plant cell wall, the most abundant polysaccharide on Earth, and an important renewable resource. Basidiomycetous fungi are among its most potent degraders because many species grow on dead wood or litter which are rich in cellulose. For the degradation of cellulose, basidiomycetes utilize a set of hydrolytic enzymes typically composed of endoglucanase, cellobiohydrolase and β -glycosidase [225, 226, 227]. For instance, *Pleurotus ostreatus* produces the cellulolytic and hemicellulolytic enzymes endo-1,4- β -glucanase, exo-1,4- β -glucanase, 1,4- β -glucosidase, endo-1,4- β -xylanase, 1,4- β -xylosidase, endo-1,4- β -mannanase and 1,4- β -mannosidase and ligninolytic enzymes Mn-peroxidase and laccase during growth on wheat straw [228].

Lignin is a complex chemical compound most commonly derived from wood, and an integral part of the secondary cell walls of plants and some algae. Several fungi have been studied for the biodegradation of lignin. Thus, agro-industrial wastes containing lignocellulose can be upgraded by solid state fermentation [229] other than biopulping during which the selective conversion of lignin is required. Several fungi (e.g. *Pleurotus* sp., *Schizophyllum* sp., *Trametes versicolor*, *Lentinus crinitus*, *Aspergillus fumigatus*, *Stemphylium vermiculosum*, *Paecilolomices carneus*, *Ceriporiopsis subvermisporea*, and species of the genus *Phlebia*) were able to grow on different agro-industrial wastes, obtaining high biodelignification [230-237]. Moreover, fungi belonging to genus *Aspergillus*, *Trichoderma*, *Phanerochaete* and *Coprinus* are known to decompose paddy straw, corn straw, wheat straw and horticultural wastes, whereas *Pleurotus sajor-caju*, *P. platypus* and *P. citrinopileatus* are known to colonize coir fibre, cotton stalks and sorghum stover. These fungi may be specific for each substrate and can be used as an effective tool for in situ degradation of lignin residues [238-243].

Trametes versicolor contributed to improving the biodegradability of Norway spruce chips from the paper industry [244] and gave better results in the removal of poplar chips than the other fungi tested, *Phanaerochaete chrysosporium* and *Pycnoporus sanguineus* [245]. Thus, Elissetche *et al.* [246] studied the biodegradation of *Drimys winteri* and *Nothofagus dombeyi*, two native Chilean wood species, by *Ganoderma australe*, which is responsible for a unique field biodegradation process resulting in completely white-rotted logs known as "palo podrido" in southern Chile.

Nyochenbeng *et al.* [247, 248] studied edible white rot fungi for selective plant biomass transformation and recycling in a sustainable ecological advanced life support (ALS) needed for extraterrestrial expeditions, such as the mission to Mars. *Pleurotus ostreatus* ('Grey Dove'), *P. pulmonarius*, *P. eryngii*, and four shiitake mushroom (*Lentinula edodes*) strains were used in the study on processed residues.

Selective degradation of lignocellulose by bamboo white rot fungi was initially studied. Zhang *et al.* [249, 250] investigated the degradation of bamboo residues by *Coriolus versicolor* B1 and *Trametes* spp. B1, having apparent degradation selectivity for hemicellulose and lignin. *Echinodontium taxodii* 2538 and *Trametes versicolor* G20 were selected for the biological pretreatment of bamboo culms (*Phyllostachys pubescens*), increasing the sugar yield of bamboo culms [251]. White-rot fungi were also capable of degrading the effluent from *Eucalyptus* chemithermomechanical pulp (CTMP) [252] and david poplar wood living on broad-leaf trees. For instance, *Funalia gallica*, *Lenzites tricolor*, *Phellinus igniarius*, *Polyporellus brumalis*, *Pseudotrametes gibbosa*, and *Pycnoporus sanguineus* reduced phenolic acids in primitive david poplar wood and wood degradation [253].

A strain of non white-rot fungi isolated from soil, *Penicillium simplicissimum*, showed different ligninolytic ability from white-rot fungi. The lignin degradation by *P. simplicissimum* happened mainly during the primary metabolism and it was greatly influenced by the pH of the media, the concentration of Cu^{2+} and Mn^{2+} [254]. *P. simplicissimum* was also tested with *Aspergillus niger* to test its capacity to decompose hydroxybenzene and nonhydroxybenzene lignin compounds of low molecular weight. Five different enzymes, lignin peroxidase, manganese peroxidase, laccase, cellulase and hemicellulase, were believed to be the most important catalysts in biodegrading process, and they always worked synergistically [255].

Lastly, biodegradation by brown-rot fungi is quantitatively one of the most important fates of lignocellulose in nature. *Gloeophyllum trabeum* and *Fomitopsis* sp. IMER2 were investigated for the biodegradation of different samples of lignin. *G. trabeum* resulted in a marked, non-selective depletion of all intermonomer side-chain linkages in the lignin [256], and *Fomitopsis* sp. IMER2 was used for the treatment of black liquor by biological acidification for the pre-

cipitation of alkali lignin [257]. Furthermore, *Piptoporus betulinus*, a common wood-rotting fungus parasitic for birch (*Betula* species), was able to degrade the lignin of birch wood [258, 259].

Polymers

The increasing consumption of plastics has generated environmental problems because it takes more than a hundred years for a discarded polymer to degrade. The ideal plastic should present desirable industrial properties and be degradable within a satisfactory time period [260].

The biodegradation ability of fungi is being investigated for polymers and plastics [261]. One example is the biodegradation of plasticized polyvinyl chloride (pPVC) by *Penicillium janthinellum* and *Doratomyces* spp. in grassland soil. The incorporation of biocides into pPVC was also studied affecting both fungal growth and the richness of species isolated [262]. Moreover, *Gloeophyllum trabeum* were used to degrade poly(vinylalcohol)(PVA) films [263, 264]. Alariqi *et al.* [265] investigated the effect of sterilization on the biodegradation by *Aspergillus niger* of polyolefins which are widely used as part of biomedical devices and food packaging after sterilization.

Attending to the biodegradability of several polymers, the degradation of a blend of the copolymer poly(hydroxybutyrate-hydroxyvalerate), PHB-HV, which is a natural, biodegradable and biocompatible thermoplastic, was studied by a mixed culture of *Phanerochaete chrysosporium* and *Talaromyces wortmannii*. The results showed that the biodegradation of the blend was a function of time, with the appearance of terminal carboxylic groups [266]. In addition, carboxymethylchitosan-g-medium chain length polyhydroxyalkanoates polyhydroxyalkanoates (mcl-PHA) were biodegraded by *Aspergillus fumigatus* 202 with a 93% weight loss of the graft [267].

Some studies were done to elucidate the microbial communities responsible for the decomposition of poly(ϵ -caprolactone) (PCL), poly(butylene succinate) (PBS), poly(butylene succinate and adipate) (PBSA), and polylactide (PLA) [268, 269, 270]. Fungi isolated from various soil environments were investigated to biodegrade poly(butylene succinate) (PBSu), such as KTF003, KTF004, and NKCM1001 strains, which have also been reported to be P(3HB)-degraders. Mesophilic strain NKCM1003 exhibited the highest poly(alkylene succinate) (PESu) hydrolytic activity among all the isolates [271, 272]. Biodegradation of polyethylene (DPE) was reported by white-rot fungi *Phanerochaete chrysosporium* was, *Talaromyces wortmannii*, and *Penicillium frequentans* showing high degradation [273, 274, 275].

Cosgrove *et al.* [276] studied the biodegradation of polyester-polyurethane (PU) showing that *Geomyces pannorum* and a *Phoma* species were the dominant species in soil fungal communities involved in the biodegradation of PU. Another recalcitrant synthetic polymer polyamide-6, generally known as nylon-6 was tested for biodegradation by fungi: five *Fusarium* spp., two *Phanerochaete chrysosporium* strains, four *Aspergillus* spp. and *Penicillium* spp., three *Cladosporium* spp. and *Ulocladium* spp., two *Trichoderma* spp., and one strain each of *Gliocladium roseum*, *Pithomyces chartarum*, *Trichotecium roseum*, and *Mucor hiemalis*. The

study showed that only white rot fungi are able to break down nylon-6 [277]. On the other hand, the biodegradation of the aliphatic polyester resin Bionolle was carried out by the filamentous fungi *Aspergillus niger* and *Penicillium funiculosum*, and *Chlorella* sp., *Lemna minor*, *Brassica rapa*, *Daphnia magna* and *Allium cepa*. The products of hydrolytic degradation did not negatively affect the organisms living in the environment [278]. The aerobic biological degradation by fungi of the synthetic aliphatic-aromatic co-polyester Ecoflex (BASF) was also studied. Weight loss was not as obvious as visual degradation and suggested broader types of microbial attack [279].

Other Industries

Fungi are capable of mineralizing a wide variety of toxic xenobiotics due to the non-specific nature of their extracellular enzymes. For instance, anaerobically digested molasses spent wash (DMSW) is a dark-brown-coloured recalcitrant effluent which has a high chemical oxygen demand (COD) and high pollution potential. Fungi such as *Aspergillus*, *Rhizopus* and *Fusarium* were able to effectively degrade DMSW [280].

Wastes from the agricultural activities were used for biodegradation. The biodegradation of untreated fertilizer industry effluent using native fungus *Aspergillus niger* and non-native, white-rot fungus *Phanerochaete chrysosporium* was studied [281].






Kalyankar *et al.* [282] investigated the degradation of India dyes by fungi from the poultry industry in Maharashtra State. Of the fungi isolated from the soil of poultry farms, *Chrysosporium tropicum*, *C. keratinophilum*, *Microsporium cannis*, *Trichophyton verrucosum*, and *T. equinum* were found to be the most dominant. *Aspergillus* sp. was the most efficient microorganism in removing ammonia from the natural sources-poultry farm and agricultural fields [283].

Penicillium sp. P6 was isolated from coal mine soil at the Qiantong colliery, Liaoning Province, North-west China, as was able to degrade Chinese lignite effectively [284]. Deuteromycete *Neosartorya fischeri*, degraded coal in the Witbank coal mining area of South Africa [285].

Biodegradation of sugar industry wastewater using the fungi *Aspergillus niger* and *Phanerochaete chrysosporium* is an effective pollution abatement solution for wastewater treatment. The fungus can degrade 98.92 % of COD and 99.86 % of biochemical oxygen demand (BOD₅) in 168 h of incubation at optimum biological process conditions. The color removal of the effluents is 99.34 % at optimum incubation time with optimum bioprocess parameters [286, 287].

One of the foremost environmental concerns in developing countries today is Solid Waste Management. In India, degradation of fruit waste was investigated by aerobic composting [288]. Garbage biodegradation was also studied by white-rot fungi [289]. Sludge degradation and bioflocculation were studied using pellet-forming filamentous fungi isolated from municipal wastewater sludge [290].

Tannery wastewater is a powerful pollutant especially due to its high chemical oxygen demand (COD). *Trametes versicolor* was examined to remove color from secondary treated tannery wastewater giving a maximum color removal efficiency of 64% [291]. *Fusarium culmorum* and *Muscador*



albus were used to treat human and/or animal waste products with a good results [292].

Lastly, the accumulation of munitions wastes in the environment has damaged many ecosystems because of their explosive properties and these compounds are biological poisons. Biodegradation by fungi is being investigated. For instance, the biodegradation of diazodinitrophenol (DDNP) wastewater was carried out by a white rot fungus cultivated and domesticated at a laboratory *in situ*. The removal of aniline compounds and nitro compounds were over 99.9%, reaching the National First-degree Wastewater Discharge Standard [293].

Undersea deposition of unexploded ordnance (UXO) constitutes a potential source of contamination of marine environments by hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX) (**94**) and octahydro-1,3,5,7-tetranitro-1,3,5,7-tetrazine (HMX) (**95**). Using sediment from a coastal UXO field, Oahu Island, Hawaii, four novel aerobic RDX-degrading fungi HAW-OCF1, HAW-OCF2, HAW-OCF3 and HAW-OCF5 were isolated and tentatively identified as members of *Rhodotorula*, *Bullera*, *Acremonium* and *Penicillium*, respectively. The four isolates mineralized 15-34% of RDX (**94**) [294]. Royal Demolition Explosive (RDX) (**94**) was also degraded by white-rot fungi, showing that the removal efficiency in wastewater could reach 87% under optimum conditions [295, 296].

Degradation of the emerging contaminant CL-20 (2,4,6,8,10,12-hexanitro-2,4,6,8,10,12-hexaazaisowurtzitane) (**96**), was reported using *Phanerochaete chrysosporium*. Another ligninolytic fungus, *Irpex lacteus*, was also able to degrade CL-20 (**96**), but as for *P. chrysosporium*, no early intermediates were observed. The intermediate was thus tentatively identified as a doubly denitrated CL-20 (**96**) product [297].

CONCLUSIONS

Biocatalysis has been demonstrated to be a powerful tool for the pollutants degradation, which is a priority for scientists in the current industrialized world.

Chemical methods have several disadvantages in the degradation of pollutants because they usually use contaminant catalysts and their use in large-scale contaminated field sites is difficult. On the other hand, biocatalytic reactions can be carried out at ambient temperature and atmospheric pressure, under safety, health, environmental, and economical conditions. In this regard, fungi have been shown to possess a variety of catabolic activities that can be harnessed to transform contaminants in less toxic compounds for the environment.

Most of the research on fungal bioremediation has been conducted on laboratory scale and conditions, so further work is required to study these capacities taking into account the natural variables and their applicability in large-scale contaminated fields.

In addition, the screening of new fungal strains with interesting enzymatic activities is necessary for the degradation of the new pollutants from the increasing industry contamination. This microorganism screening, in combination with current biotechnologies such as genetic engineering, will pave the way to the future use of fungal whole cells and enzymes for bioremediation.

REFERENCES

- [1] Arun, A.; Raja, P.P.; Arthi, R.; Ananthi, M.; Kumar, K.S.; Eyini, M. Polycyclic aromatic hydrocarbons (PAHs) biodegradation by basidiomycetes fungi, *Pseudomonas* isolate, and their cocultures: comparative in vivo and in silico approach. *Appl. Biochem. Biotechnol.*, **2008**, *151*(2-3), 132-142.
- [2] Husaini, A.; Roslan, H.A.; Hii, K.S.Y.; Ang, C.H. Biodegradation of aliphatic hydrocarbon by indigenous fungi isolated from used

- motor oil contaminated sites. *World J. Microb. Biotechnol.*, **2008**, 24(12), 2789-2797.
- [3] Bolong, N.; Ismail, A.F.; Salim, M.R.; Matsuura, T. A review of the effects of emerging contaminants in wastewater and options for their removal. *Desalination*, **2009**, 239(1-3), 229-246.
- [4] Prenafeta-Boldu, F.X.; Summerbell, R.; Sybren H.G. Fungi growing on aromatic hydrocarbons: biotechnology's unexpected encounter with biohazard? *FEMS Microbiol. Rev.*, **2006**, 30(1), 109-30.
- [5] Hou, C.; Ma, G. Performance experiment on aerobic biodegradation of benzene, toluene and xylene by fungus *Trichoderma viride* Pers.ex Fr. *Huanjing Gongcheng*, **2007**, 25(3), 45-47.
- [6] Viguera, G.; Shirai, K.; Martins, D.; Franco, T.T.; Fleuri, L.F.; Revah, S. Toluene gas phase biofiltration by *Paecilomyces lilacinus* and isolation and identification of a hydrophobic protein produced thereof. *Appl. Microbiol. Biotechnol.*, **2008**, 80(1), 147-154.
- [7] Lee, J.W.; Lee, S.M.; Hong, E.J.; Jeung, E.B.; Kang, H.Y.; Kim, M.K.; Choi, I.G. Estrogenic reduction of styrene monomer degraded by *Phanerochaete chrysosporium* KFR1 20742. *J. Microbiol.*, **2006**, 44(2), 177-184.
- [8] Cheng, G.; Li, P. Phytoremediation and microbial remediation of petroleum contaminated soil. *Huanjing Gongcheng Xueba*, **2007**, 1(6), 91-96.
- [9] Li, S.; Li, F.; Zhang, Z.; Luo, Y.; Yang, S.; Wei, B. In-situ remediation of microorganisms in frozen and thawed petroleum-contaminated soil from Liaohe Oil Field, Liaoning Province. *Liaoning Gongcheng Jishu Daxue Xuebao*, **2008**, 27(4), 599-601.
- [10] Kasumova, S.Y.; Babaeva, I. Several physiological and biochemical properties of naphthalen petroleum fungi-destructors. *Khabarlar-Azərbaycan Milli Elmlər Akademiyası, Biologiya Elmləri*, **2007**, (5-6), 123-128.
- [11] Kasumova, S.Y. Culturing micromycetes on the medium with naphthalen petroleum. *Khabarlar - Azərbaycan Milli Elmlər Akademiyası, Biologiya Elmləri*, **2005**, (3-4), 154-160.
- [12] Li, Y.Q.; Liu, H.F.; Tian, Z.L.; Zhu, L.H.; Wu, Y.H.; Tang, H.Q. Diesel pollution biodegradation: synergetic effect of Mycobacterium and filamentous fungi. *Biomed. Environ. Sci.*, **2008**, 21(3), 181-187.
- [13] Ogbo, E.M.; Okhuoya, J.A. Biodegradation of aliphatic, aromatic, resinic and asphaltic fractions of crude oil contaminated soils by *Pleurotus tuber-regium* Fr. Singer - a white rot fungus. *Afr. J. Biotechnol.*, **2008**, 7(23), 4291-4297.
- [14] Yoshioka, T.; Komuro, M. The biodegradation of the oil using fungi isolated from polluted oils. *Kaijo Hoan Daigakko Kenkyu Hokoku, Rikogaku-kei*, **2006**, 50(1-2), 1-9.
- [15] Sanyal, P.; Samaddar, P.; Paul, A.K. Degradation of poly(3-hydroxybutyrate) and poly(3-hydroxybutyrate-co-3-hydroxyvalerate) by some soil *Aspergillus* spp. *J. Polym. Environ.*, **2006**, 14(3), 257-263.
- [16] Valentin, L.; Feijoo, G.; Moreira, M.T.; Lema, J.M. Biodegradation of polycyclic aromatic hydrocarbons in forest and salt marsh soils by white-rot fungi. *Int. Biodeterior. Biodegrad.*, **2006**, 58(1), 15-21.
- [17] Bishnoi, K.; Kumar, R.; Bishnoi, N.R. Biodegradation of polycyclic aromatic hydrocarbons by white rot fungi *Phanerochaete chrysosporium* in sterile and unsterile soil. *J. Sci. Ind. Res.*, **2008**, 67(7), 538-542.
- [18] D'Annibale, A.; Ricci, M.; Leonardi, V.; Quarantino, D.; Mincione, E.; Petruccioli, M. Degradation of aromatic hydrocarbons by white-rot fungi in a historically contaminated soil. *Biotechnol. Bioeng.*, **2005**, 90(6), 723-731.
- [19] Valentin, L.; Lu-Chau, T.A.; Lopez, C.; Feijoo, G.; Moreira, M.T.; Lema, J.M. Biodegradation of dibenzothiophene, fluoranthene, pyrene and chrysene in a soil slurry reactor by the white-rot fungus *Bjerkandera* sp. BOS55. *Process Biochem.*, **2007**, 42(4), 641-648.
- [20] Leonardi, V.; Sasek, V.; Petruccioli, M.; D'Annibale, A.; Erbanova, P.; Cajthaml, T. Bioavailability modification and fungal biodegradation of PAHs in aged industrial soils. *Int. Biodeterior. Biodegrad.*, **2007**, 60(3), 165-170.
- [21] Byss, M.; Elhottova, D.; Triska, J.; Baldrian, P. Fungal bioremediation of the creosote-contaminated soil: Influence of *Pleurotus ostreatus* and *Irpex lacteus* on polycyclic aromatic hydrocarbons removal and soil microbial community composition in the laboratory-scale study. *Chemosphere*, **2008**, 73(9), 1518-1523.
- [22] Galli, E.; Rapana, P.; Tomati, U.; Polcaro, C.M.; Brancaloni, E.; Frattoni, M. Degradation of creosote by *Pleurotus ostreatus* mycelium in creosote-treated wood. *Fresen. Environ. Bull.*, **2006**, 15(8a), 720-723.
- [23] Wang, X.; Gong, Z.; Li, P.; Zhang, L.; Hu, X. Degradation of pyrene and benzo(a)pyrene in contaminated soil by immobilized fungi. *Environ. Eng. Sci.*, **2008**, 25(5), 677-684.
- [24] D'Annibale, A.; Rosetto, F.; Leonardi, V.; Federici, F.; Petruccioli, M. Role of autochthonous filamentous fungi in bioremediation of a soil historically contaminated with aromatic hydrocarbons. *Appl. Environ. Microbiol.*, **2006**, 72(1), 28-36.
- [25] Guiraud, P.; Bonnet, J.L.; Boumendjel, A.; Kadri-Dakir, M.; Dusser, M.; Bohatier, J.; Steiman, R. Involvement of *Tetrahymena pyriformis* and selected fungi in the elimination of anthracene, and toxicity assessment of the biotransformation products. *Ecotox. Environ. Safe.*, **2008**, 69(2), 296-305.
- [26] Hadibarata, T.; Tachibana, S.; Itoh, K. Biodegradation of phenanthrene by fungi screened from nature. *Pak. J. Biol. Sci.*, **2007**, 10(15), 2535-2543.
- [27] Lisowska, K.; Bizukojc, M.; Długonski, J. An unstructured model for studies on phenanthrene bioconversion by filamentous fungus *Cunninghamella elegans*. *Enzyme Microb. Tech.*, **2006**, 39(7), 1464-1470.
- [28] Kim, J.D.; Lee, C.G. Microbial degradation of polycyclic aromatic hydrocarbons in soil by bacterium-fungus co-cultures. *Biotechnol. Bioprocess Eng.*, **2007**, 12(4), 410-416.
- [29] Chen, F.; Liang, L.; Tang, Y.; Mao, L. Research on anthracene removal from water by immobilized *Fusarium oxysporum*. *Zhongguo Jishui Paishui*, **2007**, 23(21), 77-80.
- [30] Chen, F.; Tang, Y.; Mao, L.; Liang, L. Research on biodegradation characteristics of phenanthrene by *Fusarium solani* strain. *Jiangsu Keji Daxue Xuebao, Ziran Kexueban*, **2008**, 22(3), 72-76.
- [31] Li, P.; Li, H.; Stagnitti, F.; Wang, X.; Zhang, H.; Gong, Z.; Liu, W.; Xiong, X.; Li, L.; Austin, C.; Barry, D.A. Biodegradation of pyrene and phenanthrene in soil using immobilized fungi *Fusarium* sp. *Bull. Environ. Contam. Tox.*, **2005**, 75(3), 443-450.
- [32] Li, H.; Lan, W.; Lin, Y. Biotransformation of 1,2,3,4-tetrahydronaphthalene by marine fungus *Hypoxylon oceanicum*. *Fenxi Ceshi Xuebao*, **2005**, 24(4), 45-47.
- [33] Atagana, H.I. Fungal bioremediation of PAHs in the presence of heavy metals in soil. *Proceedings of the International Conference on Remediation of Chlorinated and Recalcitrant Compounds*, United States, **2006**, a.73.ppt/1-a.73.ppt/8.
- [34] Su, D.; Li, P.; Frank, S.; Xiong, X. Biodegradation of benzo[a]pyrene in soil by *Mucor* sp. SF06 and *Bacillus* sp. SB02 co-immobilized on vermiculite. *J. Environ. Sci.*, **2006**, 18(6), 1204-1209.
- [35] Su, D.; Li, P.; Wang, X.; Stagnitti, F.; Xiong, X. Biodegradation of benzo[a]pyrene in soil by immobilized fungus. *Environ. Eng. Sci.*, **2008**, 25(8), 1181-1188.
- [36] Zang, S.; Li, P.; Zhang, Y.; Wang, J.; Xu, H. Research advances in microbial metabolic pathway of benzo[a]pyrene in contaminated soils. *Shengtaixue Zazhi*, **2006**, 25(8), 978-982.
- [37] Chulalaksananukul, S.; Gadd, G. M.; Sangvanich, P.; Sihanonth, P.; Piapukiew, J.; Vangnai, A.S. Biodegradation of benzo(a)pyrene by a newly isolated *Fusarium* sp. *FEMS Microbiol. Lett.*, **2006**, 262(1), 99-106.
- [38] Su, D.; Li, P.; Ju, J.L. Degradation of pyrene and benzo- α -pyrene in soil by six strains of fungi and its kinetics. *Zhongguo Huanjing Kexue*, **2006**, 26(2), 188-191.
- [39] Cajthaml, T.; Erbanova, P.; Sasek, V.; Moeder, M. Breakdown products on metabolic pathway of degradation of benzo[a]anthracene by a ligninolytic fungus. *Chemosphere*, **2006**, 64(4), 560-564.
- [40] Zang, S.; Li, P.; Li, W.; Zhang, D.; Hamilton, A. Degradation mechanisms of benzo[a]pyrene and its accumulated metabolites by biodegradation combined with chemical oxidation. *Chemosphere*, **2007**, 67(7), 1368-1374.
- [41] Li, J.; Zhang, Z. Application of fungi in treatment of phenol-containing wastewater. *Huanjing Gongcheng Xuebao*, **2007**, 1(2), 20-24.
- [42] Strong, P.J.; Burgess, J.E. Treatment methods for wine-related and distillery wastewaters: A review. *Bioremediation J.*, **2008**, 12(2), 70-87.

- [43] Tripathi, A.K.; Harsh, N.S.K.; Gupta, N. Fungal treatment of industrial effluents: a mini-review. *Life Sci. J.*, **2007**, 4(2), 78-81.
- [44] Ren, D.; Yan, K.; Zhang, X. Study on effects of co-substrates on the biodegradation of quinoline by white rot fungi. *Huanjing Wuran Yu Fangzhi*, **2008**, 30(8), 32-35.
- [45] D'Annibale, A.; Quarantino, D.; Federici, F.; Fenice, M. Effect of agitation and aeration on the reduction of pollutant load of olive mill wastewater by the white-rot fungus *Panus tigrinus*. *Biochem. Eng. J.*, **2006**, 29(3), 243-249.
- [46] Sampedro, I.; D'Annibale, A.; Ocampo, J.A.; Stazi, S.R.; Garcia-Romera, I. Solid-state cultures of *Fusarium oxysporum* transform aromatic components of olive-mill dry residue and reduce its phytotoxicity. *Bioresour. Technol.*, **2007**, 98(18), 3547-3554.
- [47] Cai, W.; Li, J.; Zhang, Z. The characteristics and mechanisms of phenol biodegradation by *Fusarium* sp. *J. Hazard. Mater.*, **2007**, 148(1-2), 38-42.
- [48] Li, J.; Zhang, Z. Degradation capability of *Fusarium* sp. HJ01 on phenol. *Huagong Huanbao*, **2006**, 26(5), 353-356.
- [49] Prabu, P.C.; Udayasoorian, C. Biodecolorization of phenolic paper mill effluent by ligninolytic fungus *Trametes versicolor*. *J. Biol. Sci.*, **2005**, 5(5), 558-561.
- [50] Udayasoorian, C.; Prabu, P.C. Biodegradation of phenols by ligninolytic fungus *Trametes versicolor*. *J. Biol. Sci.*, **2005**, 5(6), 824-827.
- [51] Yemendzhiev, H.; Gerginova, M.; Krastanov, A.; Stoilova, I.; Alexieva, Z. Growth of *Trametes versicolor* on phenol. *J. Ind. Microbiol. Biotechnol.*, **2008**, 35(11), 1309-1312.
- [52] Cheng, G.; Sun, H.; Du, W. Research progress in degradation of organic contaminants by mycorrhizal fungi. *Turang Tongbao*, **2007**, 38(4), 799-803.
- [53] Zeng, R.; Mallik, A.U. Selected ectomycorrhizal fungi of Black Spruce (*Picea mariana*) can detoxify phenolic compounds of *Kalmia angustifolia*. *J. Chem. Ecol.*, **2006**, 32(7), 1473-1489.
- [54] Yin, P.; Li, P. Recent advances in bioremediation mechanism of PCBs contamination in soil and sediment. *Huanjing Wuran Zhili Jishu Yu Shebei*, **2005**, 6(1), 1-7.
- [55] Stoilova, I.S.; Iemendzhieva, H.T.; Krastanov, A.I. Effect of the inoculum age of *Aspergillus awamori* NRRL 3112 conidia on biodegradation of phenolic compounds and their mixtures. *Nauchni Trudove - Universitet po Khraanitelni Tekhnologii, Plovdiv*, **2007**, 54(1), 371-376.
- [56] Stoilova, I.; Krastanov, A.; Yemendzhiev, H.; Alexieva, Z. Influence of concentration of conidia of *Aspergillus awamori* NRRL 3112 on the catabolism of aromatic hydrocarbons and their mixtures. *EJEAFChe, Electronic J. Environ., Agric. Food Chem.*, **2008**, 7(4), 2833-2843.
- [57] Stoilova, I.; Krastanov, A.; Yanakieva, I.; Kratchanova, M.; Yemendzhiev, H. Biodegradation of mixed phenolic compounds by *Aspergillus awamori* NRRL 3112. *Int. Biodeterior. Biodegrad.*, **2007**, 60(4), 342-346.
- [58] Farrell, R.L.; Lamar, R.T.; White, R.B. An isolated culture of a fungal isolate that is effective to degrade hydrocarbons, and bioremediation uses. *N.Z.*, **2008**, 54pp.
- [59] Kamei, I.; Sonoki, S.; Haraguchi, K.; Kondo, R. Fungal bioconversion of toxic polychlorinated biphenyls by white-rot fungus *Phlebia brevispora*. *Appl. Microbiol. Biotechnol.*, **2006**, 73(4), 932-940.
- [60] Kamei, I.; Kogura, R.; Kondo, R. Metabolism of 4,4'-dichlorobiphenyl by white-rot fungi *Phanerochaete chrysosporium* and *Phanerochaete* sp. MZ142. *Appl. Microbiol. Biotechnol.*, **2006**, 72(3), 566-575.
- [61] Peng, D.; Zeng, G.M.; Chen, Y.N.; Yu, M.; Hu, S. Research progress on degradation of chlorophenols by using *Phanerochaete chrysosporium* biotechnology. *Shengtaixue Zazhi*, **2007**, 26(10), 1657-1664.
- [62] Gusse, A.C.; Miller, P.D.; Volk, T.J. White-rot fungi demonstrate first biodegradation of phenolic resin. *Environ. Sci. Technol.*, **2006**, 40(13), 4196-9.
- [63] Bath, H.K.; Arora, D.S. Remediation of chlorobenzoic acids by some white-rot fungi. *Proceedings of the International Conference on Remediation of Chlorinated and Recalcitrant Compounds*, United States, **2006**, a.46.ppr/1-a.46.ppr/9.
- [64] Nakagawa, A.; Osawa, S.; Hirata, T.; Yamagishi, Y.; Hosoda, J.; Horikoshi, T. 2,4-Dichlorophenol degradation by the soil fungus *Mortierella* sp. *Biosci. Biotechnol. Biochem.*, **2006**, 70(2), 525-527.
- [65] Jiang, X.; Zeng, G.; Huang, D.; Chen, Y.; Chen, X.; Huang, G. Remediation of pentachlorophenol-contaminated soil by composting with inoculation of white rot fungi. *Huanjing Kexue*, **2006**, 27(12), 2553-2557.
- [66] Rubilar, O.; Feijoo, G.; Diez, C.; Lu-Chau, T.A.; Moreira, M.T.; Lema, J.M. Biodegradation of pentachlorophenol in soil slurry cultures by *Bjerkandera adusta* and *Anthracophyllum discolor*. *Ind. Eng. Chem. Res.*, **2007**, 46(21), 6744-6751.
- [67] Zeng, B.; Ning, D.L.; Wang, H. Preliminary study on biodegradation of pentachlorophenol by white-rot fungus. *Huanjing Huaxue*, **2008**, 27(2), 181-185.
- [68] Ford, C.I.; Walter, M.; Northcott, G.L.; Di, Hong J.; Cameron, K.C.; Trower, T. Fungal inoculum properties: Extracellular enzyme expression and pentachlorophenol removal by New Zealand *Trametes* species in contaminated field soils. *J. Environ. Quality*, **2007**, 36(6), 1749-1759.
- [69] Huang, Y.; Yang, Q.; Ao, X. Tolerance and physiological responses of ectomycorrhizal fungi to pentachlorophenol. *Huanjing Kexue Xuebao*, **2008**, 28(10), 2078-2083.
- [70] Machado, K.M.G.; Matheus, D.R.; Monteiro, R.T.R.; Bononi, V.L.R.. Biodegradation of pentachlorophenol by tropical basidiomycetes in soils contaminated with industrial residues. *World J. Microbiol. Biotechnol.*, **2005**, 21(3), 297-301.
- [71] Taseli, B.K.; Gokcay, C.F. Degradation of chlorinated compounds by *Penicillium camemberti* in batch and up-flow column reactors. *Process Biochem.*, **2005**, 40(2), 917-923.
- [72] Junghanns, C.; Moeder, M.; Krauss, G.; Martin, C.; Schlosser, D. Degradation of the xenoestrogen nonylphenol by aquatic fungi and their laccases. *Microbiology*, **2005**, 151(1), 45-57.
- [73] Soares, A.; Jonasson, K.; Terrazas, E.; Guieysse, B.; Mattiasson, B. The ability of white-rot fungi to degrade the endocrine-disrupting compound nonylphenol. *Appl. Microbiol. Biotechnol.*, **2005**, 66(6), 719-725.
- [74] Shin, E.H.; Choi, H.T.; Song, H.G. Biodegradation of endocrine-disrupting bisphenol A by white rot fungus *Irpex lacteus*. *J. Microbiol. Biotechnol.*, **2007**, 17(7), 1147-1151.
- [75] Lee, S.M.; Koo, B.W.; Choi, J.W.; Choi, D.H.; An, B.S.; Jeung, E.B.; Choi, I.G. Degradation of bisphenol A by white rot fungi, *Stereum hirsutum* and *Heterobasidium insulare*, and reduction of its estrogenic activity. *Biol. Pharm. Bull.*, **2005**, 28(2), 201-207.
- [76] Lee, S.S.; Choi, D.H.; Yeo, W.H.; Choi, I.G.; Jeung, E.B. Method for biological degradation of alkylphenol causing environmental pollution using specific wood degradable fungi having superior degradability on alkylphenol, particularly 4-t-octylphenol. *Repub. Korean Kongkae Taeho Kongbo*, **2005**.
- [77] van Beek, T.A.; Claassen, F.W.; Dorado, J.; Godejohann, M.; Sierra-Alvarez, R.; Wijnberg, J.B.P.A. Fungal biotransformation products of dehydroabietic acid. *J. Nat. Prod.*, **2007**, 70(2), 154-159.
- [78] Khalaf, M.A.; Meleigy, S.A. Reduction of free gossypol levels in cottonseed meal by microbial treatment. *Int. J. Agric. Biol.*, **2008**, 10(2), 185-190.
- [79] Kondo, R.; Kamei, I. Biodegradation of dioxins and PCBs by mushroom fungi. *Bio Industry*, **2006**, 23(12), 68-76.
- [80] Chang, Y. Recent developments in microbial biotransformation and biodegradation of dioxins. *J. Mol. Microbiol. Biotechnol.*, **2008**, 15(2-3), 152-171.
- [81] Xu, X.; Lin, L.; He, B.; Jiang, L.; Ye, J.; Aorigele; SunRuncang, R. Dioxins biodegradation in the bagasse solid medium and biodegradation pathway of 2,7-DICDD in the liquid media by white rot fungi. *New Technologies in Non-Wood Fiber Pulping and Paper-making*, Guangzhou, China, **2006**, 406-411.
- [82] Xu, X.; Lin, L.; He, B.; Jiang, L.; Ye, J.; Aorigele; Sun, R. Degradation of three kinds of dioxins by white rot fungi. *Yingyong Yu Huanjing Shengwu Xuebao*, **2006**, 12(5), 701-705.
- [83] Kamei, I.; Suhara, H.; Kondo, R. Phylogenetical approach to isolation of white-rot fungi capable of degrading polychlorinated dibenzo-*p*-dioxin. *Appl. Microbiol. Biotechnol.*, **2005**, 69, 358-366.
- [84] Kamei, I.; Kondo, R. Biotransformation of dichloro-, trichloro-, and tetrachlorodibenzo-*p*-dioxin by the white-rot fungus *Phlebia lindneri*. *Appl. Microbiol. Biotechnol.*, **2005**, 68(4), 560-566.
- [85] Nam, I.; Kim, Y.; Murugesan, K.; Jeon, J.; Chang, Y.; Chang, Y. Bioremediation of PCDD/Fs-contaminated municipal solid waste incinerator fly ash by a potent microbial biocatalyst. *J. Hazard. Mater.*, **2008**, 157(1), 114-121.

- [86] Yamaguchi, M.; Kamei, I.; Nakamura, M.; Takano, M.; Sekiya, A. Selection of *Pleurotus pulmonarius* from domestic basidiomycetous fungi for biodegradation of chlorinated dioxins as environmentally persistent organopollutants. *Shinrin Sogo Kenkyusho Kenkyu Hokoku*, **2007**, 6(4), 231-237.
- [87] Tachibana, S.; Kiyota, Y.; Koga, M. Bioremediation of dioxine-contaminated soil by fungi screened from nature. *Pakistan J. Biol. Sci.*, **2007**, 10(3), 486-491.
- [88] Miyoshi, S.; Kimura, K.; Matsumoto, R.; Itoh, K.; Tachibana, S. Biodegradation of 2,8-dichlorodibenzo-*p*-dioxin by fungi screened from nature. *Pakistan J. Biol. Sci.*, **2005**, 8(9), 1265-1271.
- [89] Tachibana, S.; Kiyota, Y.; Koga, M. Bioremediation of 2,3,7,8-tetrachlorodibenzo-*p*-dioxin in soil by fungi screened from nature. *Pakistan J. Biol. Sci.*, **2006**, 9(2), 217-222.
- [90] Tachibana, S.; Kiyota, Y.; Koga, M. Biodegradation of 2,7-dibenzo-*p*-dioxin and 2,4,8-trichlorodibenzofuran in soil by fungi screened from nature. *Pakistan J. Biol. Sci.*, **2005**, 8, 1751-1757.
- [91] Zhang, X.; Yan, K.; Ren, D.; Wang, H. Studies on quinoline biodegradation by a white rot fungus (*Pleurotus ostreatus* BP) in liquid and solid state substrates. *Fresen. Environ. Bull.*, **2007**, 16, 632-638.
- [92] Ignatenko, S.V.; Voloshina, I.M.; Pirog, T.P. Biodegradation of surfactants of *Rhodococcus erythropolis* EK-1 and selection of biocides for prevention of this process. *Kharchova Promislovist*, **2007**, 5 30-33.
- [93] Ventura, J.; Belmares, R.; Aguilera-Carbo, A.; Gutierrez-Sanchez, G.; Rodriguez-Herrera, R.; Aguilar, C.N. Fungal biodegradation of tannins from creosote bush (*Larrea tridentata*) and tar bush (*Flourensia cernua*) for gallic and ellagic acid production. *Food Technol. Biotechnol.*, **2008**, 46(2), 213-217.
- [94] Martin, C.; Moeder, M.; Daniel, X.; Krauss, G.; Schlosser, D. Biotransformation of the polycyclic musks HHCB and AHTN and metabolite formation by fungi occurring in freshwater environments. *Environ. Sci. Technol.*, **2007**, 41(15), 5395-5402.
- [95] Lee, S.M.; Lee, J.W.; Koo, B.W.; Kim, M.K.; Choi, D.H.; Choi, I.G. Dibutyl phthalate biodegradation by the white rot fungus, *Polyporus brumalis*. *Biotechnol. Bioeng.*, **2007**, 97(6), 1516-1522.
- [96] Wengel, M.; Kothe, E.; Schmidt, C.M.; Heide, K.; Gleixner, G. Degradation of organic matter from black shales and charcoal by the wood-rotting fungus *Schizophyllum commune* and release of DOC and heavy metals in the aqueous phase. *Sci. Total Environ.*, **2006**, 367(1), 383-393.
- [97] Ji, Y.; Wang, Q.; Huang, L.; Li, X. Application of white rot fungi in biotreatment of refractory organic pollutant. *Nongye Gongcheng Xuebao*, **2006**, 22, 211-214.
- [98] Katapodis, P.; Moukoulis, M.; Christakopoulos, P. Biodegradation of indole at high concentration by persolvent fermentation with the thermophilic fungus *Sporotrichum thermophile*. *Int. Biodeterior. Biodegrad.*, **2007**, 60(4), 267-272.
- [99] Ren, D.; Zhang, X.; Yan, K.; Yuan, S.; Lu, X. Studies on the degradation of indole using white rot fungus. *Fresen. Environ. Bull.*, **2006**, 15(10), 1238-1243.
- [100] Singh, R.K.; Kumar, S.; Kumar, S.; Kumar, A. Biodegradation kinetic studies for the removal of p-cresol from wastewater using *Gliomastix indicus* MTCC 3869. *Biochem. Eng. J.*, **2008**, 40(2), 293-303.
- [101] Elshafie, A.; AlKindi, A.Y.; Al-Busaidi, S.; Bakheit, C.; Albahry, S.N. Biodegradation of crude oil and n-alkanes by fungi isolated from Oman. *Marine Poll. Bull.*, **2007**, 54(11), 1692-1696.
- [102] Hadibarata, T.; Tachibana, S.; Itoh, K. Biodegradation of *n*-eicosane by fungi screened from nature. *Pakistan J. Biol. Sci.*, **2007**, 10, 1804-1810.
- [103] Zabielska-Matejuk, J.; Czarczyk, K. Biodegradation of new quaternary ammonium compounds in treated wood by mould fungi. *Wood Sci. Technol.*, **2006**, 40(6), 461-475.
- [104] Liang, Z.; Han, B.; Zhang, X. Research on white-rot fungus biodegradation characters of carbon tetrachloride. *Wuhan Ligong Daxue Xuebao*, **2008**, 30, 74-78, 86.
- [105] Marco-Urrea, E.; Gabarrell, X.; Sarra, M.; Caminal, G.; Vicent, T.; Reddy, C.A. Novel aerobic perchloroethylene degradation by the white-rot fungus *Trametes versicolor*. *Environ. Sci. Technol.*, **2006**, 40(24), 7796-7802.
- [106] Marco-Urrea, E.; Caminal, G.; Gabarrell, X.; Vicent, T.; Reddy, C.A. Aerobic degradation/mineralization of trichloroethylene and perchloroethylene by white-rot fungi. *Proceedings of the International In Situ and On-Site Bioremediation Symposium*, United States, **2007**, H44/1-H44/6.
- [107] Quintero, J.C.; Lu-Chau, T.A.; Moreira, M.T.; Feijoo, G.; Lema, J.M. Bioremediation of HCH present in soil by the white-rot fungus *Bjerkandera adusta* in a slurry batch bioreactor. *Int. Biodeterior. Biodegrad.*, **2007**, 60(4), 319-326.
- [108] Daoubi, M.; Duran-Patron, R.; Hernandez-Galan, R.; Benharref, A.; Hanson, James R.; Collado, I.G. The role of botrydiediol in the biodegradation of the sesquiterpenoid phytotoxin botrydial by *Botrytis cinerea*. *Tetrahedron*, **2006**, 62(35), 8256-8261.
- [109] Wang, L.; Wang, M.; Wen, Z.; Liu, Z. Studies on biotransformation identity of digitoxin by three fungi. *Tianjin Keji Daxue Xuebao*, **2008**, 23, 8-12.
- [110] Kjeldsen, P. Behaviour of cyanides in soil and groundwater: a review. *Water, Air, Soil Poll.*, **1999**, 115(1-4), 279-307.
- [111] Baxter, J.; Cummings, S.P. The current and future applications of microorganism in the bioremediation of cyanide contamination. *Antonie van Leeuwenhoek*, **2006**, 90(1), 1-17.
- [112] Ji, J.; Wang, X.; Li, F.; Zeng, Y.; Qiao, Y. Isolation screen and application of highly-effective cyanide-degrading fungus. *Gongye Shuichuli*, **2005**, 25(7), 35-38.
- [113] Zhou, X.; Liu, L.; Chen, Y.; Xu, S.; Chen, J. Efficient biodegradation of cyanide and ferrocyanide by Na-alginate beads immobilized with fungal cells of *Trichoderma koningii*. *Can. J. Microbiol.*, **2007**, 53(9), 1033-1037.
- [114] Hossain, S.M.; Das, M.; Begum, K.M.M.S.; Anantharaman, N. Studies on biodegradation of cyanide (AgCN) using *Phanerochaete chrysosporium*. *J. Institution Engineers*, **2005**, 85, 45-49.
- [115] Romero, M.C.; Hammer, E.; Hanschke, R.; Arambarri, A.M.; Schauer, F. Biotransformation of biphenyl by the filamentous fungus *Talaromyces helicus*. *World J. Microbiol. Biotechnol.*, **2005**, 21(2), 101-106.
- [116] Monrroy, M.; Baeza, J.; Freer, J.; Rodriguez, J. Degradation of tribromophenol by wood-decaying fungi and the 1,2-dihydroxybenzene-assisted Fenton reaction. *Bioremediation J.*, **2007**, 11(4), 195-200.
- [117] Monrroy, M.; Freer, J.; Baeza, J.; Rodriguez, J. Degradation of tribromophenol by wood-rot fungi and Hamilton system. *Electronic J. Biotechnol.*, **2006**, 9(3), 253-257.
- [118] Nunez, Y.O.; Salabarría, I.S.; Collado, I.G.; Hernandez-Galan, R. The antifungal activity of widdrol and its biotransformation by *Colletotrichum gloeosporioides* (penz.) Penz. & Sacc. and *Botrytis cinerea* Pers.: Fr. *J. Agric. Food Chem.*, **2006**, 54(20), 7517-7521.
- [119] Bernat, P.; Dlugonski, J. Tributyltin chloride interactions with fatty acids composition and degradation ability of the filamentous fungus *Cunninghamella elegans*. *Int. Biodeterior. Biodegrad.*, **2007**, 60(3), 133-136.
- [120] Bernat, P.; Dlugonski, J. Transformation of tributyltin (TBT) by the microscopic fungus *Cunninghamella elegans* in the presence of NaCl. *Inzynieria i Aparatura Chemiczna*, **2005**, 44, 7-8.
- [121] Hussain, S.; Arshad, M.; Saleem, M.; Khalid, A. Biodegradation of α and β -endosulfan by soil bacteria. *Biodegradation*, **2007**, 18(6), 731-740.
- [122] Bhalerao, T.S.; Puranik, P.R. Biodegradation of organochlorine pesticide, endosulfan, by a fungal soil isolate, *Aspergillus niger*. *Int. Biodeterior. Biodegrad.*, **2007**, 59(4), 315-321.
- [123] Romero, M.L.; Terrazas, E.; Bavel, B.; Mattiasson, B. Degradation of toxaphene by *Bjerkandera* sp. strain BOL13 using waste biomass as a cosubstrate. *Appl. Microbiol. Biotechnol.*, **2006**, 71(4), 549-554.
- [124] Phillips, T.M.; Seech, A.G.; Lee, H.; Trevors, J.T. Biodegradation of hexachlorocyclohexane (HCH) by microorganism. *Biodegradation*, **2005**, 16(4), 363-392.
- [125] Rigas, F.; Papadopoulou, K.; Dritsa, V.; Doulia, D. Bioremediation of a soil contaminated by lindane utilizing the fungus *Ganoderma australe* via response surface methodology. *J. Hazard. Mater.*, **2007**, 140(1-2), 325-332.
- [126] Nagpal, V.; Srinivasan, M.C.; Paknikar, K.M. Biodegradation of γ -hexachlorocyclohexane (Lindane) by a non-white rot fungus *Conidiobolus* 03-1-56 isolated from litter. *Indian J. Microbiol.*, **2008**, 48(1), 134-141.
- [127] Rigas, F.; Dritsa, V.; Marchant, R.; Papadopoulou, K.; Avramides, E.J.; Hatzianestis, I. Biodegradation of lindane by *Pleurotus ostreatus* via central composite design. *Environ. Int.*, **2005**, 31(2), 191-196.

- [128] Mohapatra, S.; Pandey, M.; Ahuja, A.K.; Veena, S.S.; Sandhya, R. Degradation of lindane and imidacloprid in soil by *Calocybe indica*. *Pestic. Res. J.*, **2008**, 20(1), 143-145.
- [129] Slaoui, M.; Ouhssine, M.; Berny, E.; Elyachoui, M. Biodegradation of carbofuran by a fungus isolated from treated soil. *Phys. Chem. News*, **2006**, 32, 116-122.
- [130] Seo, J.; Jeon, J.; Kim, S.D.; Kang, S.; Han, J.; Hur, H.G. Fungal biodegradation of carbofuran and carbofuran phenol by the fungus *Mucor ramannianus*: identification of metabolites. *Water Sci. Technol.*, **2007**, 55, 163-167.
- [131] Hossain, S.K.M.; Anantharaman, N. Studies on aerobic biodegradation of DDT using *Phanerochaete chrysosporium*. *Indian J. Environ. Protection*, **2005**, 25(5), 454-457.
- [132] Huang, Y.; Zhao, X.; Luan, S. Uptake and biodegradation of DDT by 4 ectomycorrhizal fungi. *Sci. Total Environ.*, **2007**, 385(1-3), 235-241.
- [133] Purnomo, A.S.; Kamei, I.; Kondo, R. Degradation of 1,1,1-trichloro-2,2-bis (4-chlorophenyl) ethane (DDT) by brown-rot fungi. *J. Biosci. Bioeng.*, **2008**, 105(6), 614-621.
- [134] Mukherjee, I.; Mittal, A. Dissipation of β -cyfluthrin by two fungi *Aspergillus nidulans* var. *dentatus* and *Sepedonium maheswarium*. *Toxicol. Environ. Chem.*, **2007**, 89(2), 319-326.
- [135] Seo, J.; Lee, Y.G.; Kim, S.D.; Cha, C.J.; Ahn, J.H.; Hur, H.G. Biodegradation of the insecticide N,N-diethyl-m-toluamide by fungi: Identification and toxicity of metabolites. *Arch. Environ. Contamination Toxicol.*, **2005**, 48(3), 323-328.
- [136] Ai, T.; Wang, H.; Wen, X.; Zhang, S.; Huang, J. Microbial remediation of dimethoate-contaminated soil. *Nongye Huanjing Kexue Xuebao*, **2006**, 25(5), 1250-1254.
- [137] Lee, S.; Lee, J.; Park, K.; Hong, E.; Jeung, E.; Kim, M.; Kang, H.; Choi, I. Biodegradation of Methoxychlor and its metabolites by the white rot fungus *Stereum hirsutum* related to the inactivation of estrogenic activity. *J. Environ. Sci. Health, Part B: Pestic., Food Contam., Agric. Wastes*, **2006**, 41(4), 385-397.
- [138] Osman, K.A.; Ibrahim, G.H.; Askar, A.I.; Aba Alkhail, A.R.A. Biodegradation kinetics of dicofol by selected microorganism. *Pestic. Biochem. Physiol.*, **2008**, 91(3), 180-185.
- [139] Kanaly, R.A.; Kim, I.S.; Hur, H.G. Biotransformation of 3-methyl-4-nitrophenol, a main product of the insecticide fenitrothion, by *Aspergillus niger*. *J. Agric. Food Chem.*, **2005**, 53(16), 6426-6431.
- [140] Martinez, C.O.; Silva, C.M.; Maia, A. Biodegradation of the herbicide propanil by fungi isolated from a rice rhizosphere. *Pesticidas*, **2005**, 15, 113-121.
- [141] Martinez, C.O.; Silva, C.M.; Fay, E.F.; Abakerli, R.B.; Maia, A.; Durrant, L.R. The effects of moisture and temperature on the degradation of sulfentrazone. *Geoderma*, **2008**, 147(1-2), 56-62.
- [142] Romeh, A.A. Adsorption and biodegradation of the herbicide fluometuron in liquid media. *J. Environ. Res.*, **2006**, 7, 29-47.
- [143] Ronhede, S.; Sorensen, S.R.; Jensen, B.; Aamand, J. Mineralization of hydroxylated isoproturon metabolites produced by fungi. *Soil Biol. Biochem.*, **2007**, 39(7), 1751-1758.
- [144] Kamei, I.; Kondo, R. Simultaneous degradation of commercially produce CNP herbicide and contaminated dioxin by treatment using the white-rot fungus *Phlebia brevispora*. *Chemosphere*, **2006**, 65(7), 1221-1227.
- [145] Castro, J.V., Jr.; Peralba, M.C.R.; Ayub, M.A.Z. Biodegradation of the herbicide glyphosate by filamentous fungi in platform shaker and batch bioreactor. *J. Environ. Sci. Health, Part B: Pestic., Food Contam., Agric. Wastes*, **2007**, 42(8), 883-886.
- [146] Zhu, J.; Qiao, X.; Wang, J.; Qin, S. Biodegradation of acetochlor in soil and its persistence against *Echinochloa crusgalli*. *Yingyong Shengtai Xuebao*, **2006**, 17, 489-492.
- [147] He, Y.H.; Shen, D.S.; Fang, C.R.; He, R.; Zhu, Y.M. Effects of metsulfuron-methyl on the microbial population and enzyme activities in wheat rhizosphere soil. *J. Environ. Sci. Health, Part B, Pestic., Food Contam., Agric. Wastes*, **2006**, 41(3), 269-84.
- [148] He, Y.H.; Shen, D.S.; Fang, C.R.; Zhu, Y.M. Rapid biodegradation of metsulfuron-methyl by a soil fungus in pure cultures and soil. *World J. Microbiol. Biotechnol.*, **2006**, 22(10), 1095-1104.
- [149] Bishnoi, N.R.; Garima, Fungus- an alternative for bioremediation of heavy metal containing wastewater: A review. *J. Sci. Ind. Res.*, **2005**, 64(2), 93-100.
- [150] D'Annibale, A.; Leonardi, V.; Federici, E.; Baldi, F.; Zecchini, F.; Petruccioli, M. Leaching and microbial treatment of a soil contaminated by sulphide ore ashes and aromatic hydrocarbons. *Appl. Microbiol. Biotechnol.*, **2007**, 74(5), 1135-1144.
- [151] Baldi, F.; Leonardi, V.; D'Annibale, A.; Piccolo, A.; Zecchini, F.; Petruccioli, M. Integrate approach of metal removal and bioprecipitation followed by fungal degradation of organic pollutants from contaminated soils. *Eur. J. Soil Biol.*, **2007**, 43(5-6), 380-387.
- [152] Zeng, G.; Huang, D.; Huang, G.; Hu, T.; Jiang, X.; Feng, C.; Chen, Y.; Tang, L.; Liu, H. Composting of lead-contaminated solid waste with inocula of white-rot fungus. *Bioresour. Technol.*, **2007**, 98(2), 320-6.
- [153] Huang, D.; Zeng, G.; Feng, C.; Hu, S.; Jiang, X.; Tang, L.; Su, F.; Zhang, Y.; Zeng, W.; Liu, H. Lead-contaminated lignocellulosic waste by *Phanerochaete chrysosporium* and the reduction of lead toxicity. *Environ. Sci. Technol.*, **2008**, 42(13), 4946-4951.
- [154] Akar, T.; Tunali, S.; Cabuk, A. Study on the characterization of lead(II) biosorption by fungus *Aspergillus parasiticus*. *Appl. Biochem. Biotechnol.*, **2007**, 136, 389-406.
- [155] Akar, T.; Cabuk, A.; Tunali, S.; Yamac, M. Biosorption potential of the macrofungus *Ganoderma carnosum* for removal of lead(II) ions from aqueous solutions. *J. Environ. Sci. and Health, Part A: Toxic/Hazard. Substances Environ. Eng.*, **2006**, 41(11), 2587-2606.
- [156] Tunali, S.; Akar, T.; Oezcan, A.S.; Kiran, I.; Oezcan, A. Equilibrium and kinetics of biosorption of lead(II) from aqueous solutions by *Cephalosporium aphidicola*. *Sep. Purif. Technol.*, **2006**, 47(3), 105-112.
- [157] Tunali, S.; Akar, T. Zn(II) biosorption properties of *Botrytis cinerea* biomass. *J. Hazard. Mater.*, **2006**, 131(1-3), 137-145.
- [158] Akar, T.; Tunali, S.; Kiran, I. *Botrytis cinerea* as a new fungal biosorbent for removal of Pb(II) from aqueous solutions. *Biochem. Eng. J.*, **2005**, 25(3), 227-235.
- [159] Akar, T.; Tunali, S. Biosorption characteristics of *Aspergillus flavus* biomass for removal of Pb(II) and Cu(II) ions from an aqueous solution. *Bioresour. Technol.*, **2006**, 97(15), 1780-1787.
- [160] Kiran, I.; Akar, T.; Tunali, S. Biosorption of Pb(II) and Cu(II) from aqueous solutions by pretreated biomass of *Neurospora crassa*. *Process Biochem.*, **2005**, 40(11), 3550-3558.
- [161] Bishnoi, N.R.; Kumar, R.; Bishnoi, K. Biosorption of Cr(VI) with *Trichoderma viride* immobilized fungal biomass and cell free Ca-alginate beads. *Indian J. Exp. Biol.*, **2007**, 45(7), 657-664.
- [162] Kumar, R.; Bishnoi, N.R.; Garima; Bishnoi, K. Biosorption of Chromium(VI) from aqueous solution and electroplating wastewater using fungal biomass. *Chem. Eng. J.*, **2008**, 135(3), 202-208.
- [163] Chen, G.; Zeng, G.; Tang, L.; Du, C.; Jiang, X.; Huang, G.; Liu, H.; Shen, G. Cadmium removal from simulated wastewater to biomass byproduct of *Lentinus edodes*. *Bioresour. Technol.*, **2008**, 99(15), 7034-7040.
- [164] Kelly, D.J.A.; Budd, K.; Lefebvre, D.D. The biotransformation of mercury in pH-stat cultures of microfungi. *Can. J. Bot.*, **2006**, 84(2), 254-260.
- [165] Eichlerova, I.; Homolka, L.; Nerud, F. Evaluation of synthetic dye decolorization capacity in *Ischnoderma resinosa*. *J. Ind. Microbiol. Biotechnol.*, **2006**, 33(9), 759-766.
- [166] Eichlerova, I.; Homolka, L.; Benada, O.; Kofronova, O.; Hubalek, T.; Nerud, F. Decolorization of Orange G and Remazol Brilliant Blue R by the white rot fungus *Dichomitus squalens*: Toxicological evaluation and morphological study. *Chemosphere*, **2007**, 69(5), 795-802.
- [167] Asgher, M.; Bhatti, H.N.; Ashraf, M.; Legge, R.L. Recent developments in biodegradation of industrial pollutants by white rot fungi and their enzyme system. *Biodegradation*, **2008**, 19(6), 771-783.
- [168] Hossain, S.M. Aerobic treatment of distillery wastewater using *Phanerochaete chrysosporium*. *Indian J. Environ. Protec.*, **2007**, 27(4), 362-366.
- [169] Feng, R.; Wang, H.; Zhang, X. White rot fungus biological treatment of mixed wastewater from black liquor and diosgenin. *Gongye Shuichuli*, **2006**, 26(8), 45-48.
- [170] Li, Z.X.; Zhu, J.L. Studies on immobilized white rot fungi for biodegradation of pulp liquor. *Shuichuli Jishu*, **2005**, 31(12), 23-26.
- [171] Poznyak, T.; Colindres, P.; Chairez, I. Treatment of textile industrial dyes by simple ozonation with water recirculation. *J. Mex. Chem. Soc.*, **2007**, 51(2), 81-86.
- [172] Abd-El-Rahim, W.M.; Moawad, H. Textile industry wastes, a real treat to agricultural environment in Egypt. *Environ. Sci.*, **2008**, 3(1), 134-142.
- [173] Zhang, L.F.; Sun, Y. Decolorization of dyes by fungus strain M2. *Shenyang Ligong Daxue Xuebao*, **2006**, 25(6), 83-87.

- [174] Li, X.; Lin, C.; Xu, M.; Fu, S. Screening and decolorization of synthetic dyes of a white rot fungi ZJ-6. *Jiangxi Shifan Daxue Xuebao, Ziran Kexueban*, **2006**, 30(6), 543-546.
- [175] Contreras, A.M.; Balagurusamy, N. Decolorization of Turquoise HGN by fungi isolated from contaminated soil. *Chimica Oggi*, **2007**, 25(Suppl.), 19-20.
- [176] Zhang, J.; Kan, Z.; Su, W. Study on condition of decolorization and degradation of dyes of co-culture group. *Shengwu Jishu*, **2008**, 18(1), 78-81.
- [177] Abdel-Kareem, O. Effect of selected natural dyes in reduction on colour changes of Egyptian linen textiles by fungi. *Ann. Chim.*, **2007**, 97(7), 527-540.
- [178] Kou, X.; An, L.; Zuo, Z.; Zhang, P.; Jia, J.; Shen, J. Combined white rot fungus-activated sludge process for treatment of dye wastewater. *Jingxi Huagong*, **2007**, 24(5), 500-503.
- [179] Hai, F.I.; Yamamoto, K.; Nakajima, F.; Fukushi, K. Removal of structurally different dyes in submerged membrane fungi reactor-biosorption/PAC-adsorption, membrane retention and biodegradation. *J. Membr. Sci.*, **2008**, 325(1), 395-403.
- [180] Rigas, F.; Dritsa, V. Decolourisation of a polymeric dye by selected fungal strains in liquid cultures. *Enzyme Microb. Technol.*, **2006**, 39(1), 120-124.
- [181] Eichlerová, I.; Homolka, L.; Ludmila, L.; Nerud, F. Orange G and Remazol Brilliant Blue R decolorization by white rot fungi *Dichomitus squalens*, *Ischnoderma resinosa* and *Pleurotus calyptratus*. *Chemosphere*, **2005**, 60, 398-404.
- [182] Sukumar, M.; Sivasamy, A.; Swaminathan, G. Influence of co-substrates for the decolorization of dye house effluent by *Phanerochaete chrysosporium* and *Trametes versicolor*. *Poll. Res.*, **2006**, 25(3), 557-562.
- [183] Zuo, Z.; An, L.; Mao, R.; Kou, X.; Jia, J.; Shen, J.; Zhang, P. Discolourization of dye by micro-electrolysis-P.C. process. *Jingxi Huagong*, **2007**, 24(5), 496-499.
- [184] Machado, K.M.G.; Compant, L.C.A.; Morais, R.O.; Rosa, L.H.; Santos, M.H. Biodegradation of reactive textile dyes by basidiomycetous fungi from Brazilian ecosystems *Braz. J. Microbiol.*, **2006**, 37(4), 481-487.
- [185] Zhang, X.; Yan, K.; Wang, H.; Ren, D. Study on the biodegradation mechanisms of triphenylmethane dyes by the white rot fungus in rice straw medium. *Huanjing Kexue Xuebao*, **2006**, 26(8), 1284-1289.
- [186] Tian, Y.; Yan, K.; Wang, H.; Zhang, X. Removal of malachite green and crystal violet by adsorption on rice straw and biodegradation of white rot fungus. *Huanjing Wuran Yu Fangzhi*, **2008**, 30(5), 33-37.
- [187] Zhang, X.; Yan, K.; Wang, H.; Yin, Y. Biodegradable liquid of lignocelluloses accelerate recalcitrant dye decolorization by white rot fungi. *Fresen. Environ. Bull.*, **2007**, 16(12a), 1563-1570.
- [188] Yan, K.; Tian, Y.; Wang, H.; Zhang, X. Study on adsorptive removal of dyes by white rot fungus. *Shengwu Jishu*, **2007**, 17(5), 68-71.
- [189] Malachova, K.; Pavlickova, Z.; Novotny, C.; Svobodova, K.; Lednicka, D.; Musilkova, E. Reduction in the mutagenicity of synthetic dyes by successive treatment with activated sludge and the ligninolytic fungus, *Irpex lacteus*. *Environ. Mol. Mutagen.*, **2006**, 47(7), 533-540.
- [190] Naidu, K.S.B.; Devi, K.L. Biodegradation of textile dyes by the fungus *Phanerochaete chrysosporium*. *J. Ecobiol.*, **2006**, 18(3), 277-281.
- [191] Demir, G.; Ozcan, H.K.; Tufekci, N.; Borat, M. Decolorization of Remazol Yellow RR Gran by white rot fungus *Phanerochaete chrysosporium*. *J. Environ. Biol.*, **2007**, 28(4), 813-817.
- [192] Sukumar, M.; Sivasamy, A.; Swaminathan, G.; Sai, R.C.R.; Saravanan, M. Process optimization for the decolorization of Bismarck brown by *Phanerochaete chrysosporium*. *Res. J. Biotechnol.*, **2007**, 2(3), 35-39.
- [193] Gao, D.; Wen, X.; Qian, Y. Decolorization of Reactive Brilliant Red K-2BPhy white rot fungus under sterile and non-sterile conditions. *J. Environ. Sci.*, **2006**, 18(3), 428-432.
- [194] Eichlerová, I.; Homolka, L.; Nerud, F. Synthetic dye decolorization capacity of white rot fungus *Dichomitus squalens*. *Bioresour. Technol.*, **2006**, 97, 2153-2159.
- [195] Asgher, M.; Batool, S.; Bhatti, H.N.; Noreen, R.; Rahman, S.U.; Javid Asad, M. Laccase mediated decolorization of vat dyes by *Coriolus versicolor* IBL-04. *Int. Biodeterior. Biodegrad.*, **2008**, 62(4), 465-470.
- [196] Asgher, M.; Shah, S.A.H.; Ali, M.; Legge, R.L. Decolorization of some reactive textile dyes by white rot fungi isolated in Pakistan. *World J. Microbiol. Biotechnol.*, **2006**, 22(1), 89-93.
- [197] Asgher, M.; Kausar, S.; Bhatti, H.N.; Shah, S.A.H.; Ali, M. Optimization of medium for decolorization of Solar Golden Yellow R direct textile dye by *Schizophyllum commune* IBL-06. *Int. Biodeterior. Biodegrad.*, **2008**, 61(2), 189-193.
- [198] Bhatti, H.N.; Tabassum, S.; Asgher, M. Enhanced decolourization of Solar Orange RSN by white-rot fungus *Ganoderma lucidum* IBL-05. *Asian J. Chem.*, **2008**, 20(8), 6011-6021.
- [199] Bhatti, H.N.; Akram, N.; Asgher, M. Optimization of culture conditions for enhanced decolourization of Cibacron Red FN-2BL by *Schizophyllum commune* IBL-6. *Appl. Biochem. Biotechnol.*, **2008**, 149(3), 255-264.
- [200] Machado, K.M.G.; Matheus, D.R.; Bononi, V.L.R. Ligninolytic enzymes production and Remazol Brilliant Blue R decolorization by tropical Brazilian basidiomycetes fungi. *Braz. J. Microb.*, **2005**, 36(3), 246-252.
- [201] Vitali, V.M.V.; Machado, K.M.G.; de Andrea, M.M.; Bononi, V.L.R. Screening mitosporic fungi for organochlorides degradation. *Braz. J. Microb.*, **2006**, 37(3), 256-261.
- [202] Zhao, X.; Lu, Y.; Phillips, D.R.; Hwang, H.; Hardin, I.R. Study of biodegradation products from azo dyes in fungal degradation by capillary electrophoresis/electrospray mass spectrometry. *J. Chromatogr. A*, **2007**, 1159(1-2), 217-224.
- [203] Fu, C.; Zhang, J.; Zheng, J.; Zhao, L.; Liu, J. Isolation and enhanced action of predominant fungi in biological treatment of dyeing wastewater. *Yingyong Yu Huanjing Shengwu Xuebao*, **2006**, 12(5), 693-696.
- [204] Srikanlayanukul, M.; Kitwechun, W.; Watanabe, T.; Khanongnuch, C. Decolorization of Orange II by immobilized thermotolerant white fungus *Coriolus versicolor* RC3 in packed-bed bioreactor. *Biotechnol.*, **2008**, 7(2), 280-286.
- [205] Lu, Y.; Phillips, D.R.; Lu, L.; Hardin, I.R. Determination of the degradation products of selected sulfonated phenylazonaphthol dyes treated by white rot fungus *Pleurotus ostreatus* by capillary electrophoresis coupled with electrospray ionization ion trap mass spectrometry. *J. Chromatogr. A*, **2008**, 1208(1-2), 223-231.
- [206] Lu, Y.; Hardin, I. Analysis of sulfonated azo dyes degraded by white rot fungus *Pleurotus ostreatus*. *AATCC Rev.*, **2006**, 6(1), 31-36.
- [207] Wang, L.; Hardin, I. Analysis of hydrophobic dyes' biodegradation by white rot fungus *Pleurotus ostreatus*. *Preprints of Extended Abstracts presented at the ACS National Meeting, Division of Environmental Chemistry*, **2005**, 45(2), 201.
- [208] Zhao, X.; Hardin, I.R.; Hwang, H. Biodegradation of a model azo disperse dye by the white rot fungus *Pleurotus ostreatus*. *Int. Biodeterior. Biodegrad.*, **2006**, 57(1), 1-6.
- [209] Abd El-Rahim, W.M.; Khalil, W.K.B.; Eshak, M.G. Genotoxicity studies on the removal of a direct textile dye by a fungal strain, in vivo, using micronucleus and RAPD-PCR techniques on male rats. *J. Appl. Toxicol.*, **2008**, 28(4), 484-490.
- [210] Khalaf, M.A.. Biosorption of reactive dye from textile wastewater by non-viable biomass of *Aspergillus niger* and *Spirogyra* sp. *Bioresour. Technol.*, **2008**, 99(14), 6631-6634.
- [211] Zhang, X.; Liu, Y.; Yan, K.; Wu, H. Decolorization of anthraquinone-type dye by bilirubin oxidase-producing nonligninolytic fungus *Myrothecium* sp. IMER1. *J. Biosci. Bioeng.*, **2007**, 104(2), 104-110.
- [212] Sengottuvel, R.; Srinivasan, K.; Natarajan, D.; Mohanasundari, C.; Varghese, S. Studies on textile dye degrading ability of fungi. *Asian J. Microb. Biotechnol. Environ. Sci.*, **2007**, 9(2), 277-280.
- [213] Cai, W.; Li, F.; Li, J. The study on Neutral Brilliant Blue GL decolorization and degradation characteristics by a *Fusarium* sp. HJ01 strain. *Huanjing Kexue Xuebao*, **2007**, 27(2), 213-219.
- [214] Li, J.; Li, F. Isolation of acid blue B-degrading *Fusarium* sp. HJ01 and the studies on its degradation characteristics. *Huanjing Kexue Xuebao*, **2005**, 25(12), 1641-1646.
- [215] Gupta, A.K.; Pandey, S.K. Role of fungi in bioremediation of waste water discharged from dyeing plant of a carpet industry at Varanasi. *Poll. Res.*, **2006**, 25(4), 787-791.
- [216] Yesiladali, S.K.; Pekin, G.; Bermek, H.; Arslan-Alaton, I.; Orhon, D.; Tamerler, C. Bioremediation of textile azo dyes by *Trichophyton rubrum* LSK-27. *World J. Microbiol. Biotechnol.*, **2006**, 22(10), 1027-1031.

- [217] Lakshmanan, P.; Jagadeesan, R.; Sudha, A.; Rajesh, M.; Prabhakara, S.; Prasad, M.A. Potentiality of a new mushroom fungus *Lentinus connatus* Berk. for the production of biomanure from sugarcane trash (*Saccharum officinarum* L.) and its impact on the management of groundnut root rot diseases. *Arch. Phytopathol. Plant Protec.*, **2008**, 41(4), 273-289.
- [218] Kokol, V.; Doliska, A.; Eichlerova, I.; Baldrian, P.; Nerud, F. Decolorization of textile dyes by whole cultures of *Ischnoderma resinotum* and by purified laccase and Mn-peroxidase. *Enzyme Microb. Technol.*, **2007**, 40(7), 1673-1677.
- [219] Murugesan, K.; Nam, I.; Kim, Y.; Chang, Y. Decolorization of reactive dyes by a thermostable laccase produced by *Ganoderma lucidum* in solid state culture. *Enzyme Microb. Technol.*, **2007**, 40(7), 1662-1672.
- [220] Dritsa, V.; Rigas, F.; Natsis, K.; Marchant, R. Characterization of a fungal strain isolated from a polyphenol polluted site. *Bioresour. Technol.*, **2007**, 98(9), 1741-1747.
- [221] Sukumar, M.; Sivasamy, A.; Swaminathan, G. Biological decolorization of dye house effluent by *Trametes* sp. isolated from contaminated soil. *Res. J. Biotechnol.*, **2008**, 3, 53-58.
- [222] Rajamohan, N.; Karthikeyan, C. Effect of operating parameters on aerobic treatment of dyehouse effluent by *Galactomyces geotrichum*. *Ecol. Environ. Conserv.*, **2005**, 11, 487-490.
- [223] Gavril, M.; Hodson, P.V.; McLellan, J. Decoloration of Amaranth by the white-rot fungus *Trametes versicolor*. Part I. Statistical analysis. *Can. J. Microbiol.*, **2007**, 53(2), 313-326.
- [224] Gavril, M.; Hodson, P.V. Chemical evidence for the mechanism of the biodecolorization of Amaranth by *Trametes versicolor*. *World J. Microbiol. Biotechnol.*, **2007**, 23(1), 103-124.
- [225] Baldrian, P.; Valaskova, V. Degradation of cellulose by basidiomycetous fungi. *FEMS Microbiol. Rev.*, **2008**, 32(3), 501-521.
- [226] Wilson, D. Three microbial strategies for plant cell wall degradation. *Ann. N. Y. Acad. Sci.*, **2008**, 1125, 289-297.
- [227] Wang, H.; Li, P.; Zhang, W.; Liu, G. Study on isolation of white rot fungi F8 from soil sample and its ability for degrading lignin. *Henan Shifan Daxue Xuebao, Ziran Kexueban*, **2006**, 34(2), 110-112.
- [228] Baldrian P.; Valaskova V.; Merhautova V.; Gabriel J. Degradation of lignocellulose by *Pleurotus ostreatus* in the presence of copper, manganese, lead and zinc. *Res. Microbiol.*, **2005**, 156(5-6), 670-6.
- [229] Hamza, A.S.; Darwish, G.A.M.A.; Abdel-Kawi, K.A. Bioconversion of food processing wastes by *Pleurotus ostreatus* into protein enriched product. *New Egyptian J. Microbiol.*, **2005**, 12, 240-251.
- [230] Fackler, K.; Grading, C.; Schmutzer, M.; Tavzes, C.; Burgert, I.; Schwanninger, M.; Hinterstoisser, B.; Watanabe, T.; Messner, K. Biotechnological wood modification with selective white-rot fungi and its molecular mechanism. *Food Technol. Biotechnol.*, **2007**, 45(3), 269, 276.
- [231] Liu, Z.; Wang, H.; Fang, G.; Han, S.; Li, P. Research progress of white-rot fungi application on pulp and papermaking industry. *Zhongguo Zaozhi*, **2007**, 26, 47-52.
- [232] Klyagina, Y.P.; Smirnov, V.F.; Strutchkova, I.V.; Trofimov, A.N.; Kisilitsyn, A.N. Biodegradation of particleboard lignin by microscopic fungi. *Khimiya Rastitel'nogo Syr'ya*, **2005**, 4, 41-44.
- [233] Hossain, Sk.M.; Anantharaman, N. Comparison studies on biological pulping of lignocellulosic agro-residue using *Trametes versicolor* and *Lentinus crinitus*. *Process Plant Eng.*, **2006**, 24(2), 41-46.
- [234] Hossain, Sk.M.; Anantharaman, N. Effect of glucose on lignin biodegradation of lignocellulosic agro-residue using *Lentinus crinitus*. *Bulgarian Chem. Ind.*, **2005**, 76(1-2), 50-54.
- [235] Bappuji, S.; Mohan, T.S.; Joy, B. Efficiency of lignocellulosic fungi on decomposition of coir pith. *Poll. Res.*, **2005**, 24(3), 729-733.
- [236] Watanabe, T.; Ohashi, Y.; Tanabe, T.; Honda, Y.; Messner, K. Lignin biodegradation by selective white rot fungus and its potential use in wood biomass conversion. *ACS Symp. Ser.*, **2007**, 954, 409-421.
- [237] Watanabe, T.; Ougi, T.; Nishimura, H.; Watanabe, T.; Honda, Y.; Okano, K. Free radical-mediated lignin biodegradation by selective white rot fungi and its potential use in wood biomass conversion. *Research Progress in Pulp and Papermaking*, **2006**, 864-868.
- [238] Gaind, S.; Nain, L. Chemical and biological properties of wheat soil in response to paddy straw incorporation and its biodegradation by fungal inoculants. *Biodegradation*, **2007**, 18(4), 495-503.
- [239] Gaind, S.; Lata; Goyal, D. Trace element characterization for quality evaluation of compost form amended paddy straw inoculated with fungal consortium. *Indian J. Microbiol.*, **2006**, 46(2), 127-132.
- [240] Gaind, S.; Pandey, A.K.; Lata. Biodegradation study of crop residues as affected by exogenous inorganic nitrogen and fungal inoculants. *J. Basic Microbiol.*, **2005**, 45(4), 301-311.
- [241] Hossain, Sk.M.; Anantharaman, N. Studies on biopulping of agro-residue wheat straw using *Phanerochaete chrysosporium*. *Process Plant Eng.*, **2005**, 23(3), 55-58.
- [242] Wang, H.; Du, F.; Zhang, X. Selective degradation of corn straw lignocellulose by white-rot fungi. *Huazhong Keji Daxue Xuebao, Ziran Kexueban*, **2006**, 34(3), 97-100.
- [243] Zeng, G.; Chen, Y. Screening of strains compatible with *Phanerochaete chrysosporium* and coculture for biodegradation of straw. *Changsha Ligong Daxue Xuebao, Ziran Kexueban*, **2006**, 3(3), 107-112.
- [244] van Beek, T.A.; Kuster, B.; Claassen, F.W.; Tienverier, T.; Bertaud, F.; Lenon, G.; Petit-Conil, M.; Sierra-Alvarez, R. Bio-treatment of spruce wood with *Trametes versicolor* for pitch control: Influence on extractive contents, pulping process parameters, paper quality and effluent. *Bioresour. Technol.*, **2006**, 98(2), 302-311.
- [245] Han, S.; Fang, G.; Chu, F.; Ping, L.; Sigoillot, J.C. Study on treating poplar by 4 white-rot fungi. *Research Progress in Pulp and Papermaking*, **2006**, 247-250.
- [246] Elissetche, J.; Ferraz, A.; Freer, J.; Rodriguez, J. Influence of forest on biodegradation of *Drimys winteri* by *Ganoderma australe*. *Int. Biodeterior. Biodegrad.*, **2006**, 57(3), 174-178.
- [247] Nyochembeng, L.M.; Beyl, C.A.; Pacumbaba, R.P. Nitrogen amendment enhances edible white rot fungal growth and biodegradation of containerized inedible crop residues. *Habitation*, **2006**, 11, 133-138.
- [248] Nyochembeng L.M.; Beyl C.A.; Pacumbaba R.P. Optimizing edible fungal growth and biodegradation of inedible crop residues using various cropping methods. *Bioresour. Technol.*, **2008**, 99(13), 5645-9.
- [249] Zhang, X.; Xu, C.; Wang, H. Pretreatment of bamboo residues with *Corioliolus versicolor* for enzymatic hydrolysis. *J. Biosci. Bioeng.*, **2007**, 104(2), 149-151.
- [250] Xu, C.; Wang, H.; Zhang, X.; Fu, S.; Wu, J. Lignocellulose selective degradation by white rot fungi in bamboo. *Weishengwuxue Zazhi*, **2006**, 26(2), 14-18.
- [251] Zhang, X.; Yu, H.; Huang, H.; Liu, Y. Evaluation of biological pretreatment with white rot fungi for the enzymatic hydrolysis of bamboo culms. *Int. Biodeterior. Biodegrad.*, **2007**, 60(3), 159-164.
- [252] Xu, X.; He, B.; Li, J.; Lin, L.; Chen, Y.; Li, H.; Hu, Z. Bio-discoloring of effluent from eucalyptus CTMP with white-rot fungus. *Zhongguo Zaozhi Xuebao*, **2007**, 22(3), 57-59.
- [253] Chi, Y.; Yan, H. HPLC analysis on different phenolic acids and their contents of David poplar wood degraded by 6 species of wood white-rot fungi. *Linye Kexue*, **2008**, 44(2), 116-123.
- [254] Yu, H.; Zeng, G.; Huang, G.; Huang, D.; Chen, Y. Lignin degradation by *Penicillium simplicissimum*. *Zhongguo ke xue yuan huan jing ke xue wei yuan hui "Huan jing ke xue" bian ji wei yuan hui*, **2005**, 26(2), 167-71.
- [255] Chen, F.; Xie, G.; Yu, H.; Peng, D.; Yu, M.; Huang, H.; Zeng, G. Development of lignin-biodegrading inoculant for composting. *Shengtai Yu Nongcun Huanjing Xuebao*, **2008**, 24(2), 84-87.
- [256] Yelle, D.J.; Ralph, J.; Lu, F.; Hammel, K.E. Evidence for cleavage of lignin by a brown rot basidiomycete. *Environ. Microbiol.*, **2008**, 10(7), 1844-1849.
- [257] Xiong, Z.; Zhang, X.; Wang, H.; Ma, F.; Li, L.; Li, W. Application of brown-rot basidiomycete *Fomitopsis* sp. IMER2 for biological treatment of black liquor. *J. Biosci. Bioeng.*, **2007**, 104(6), 446-450.
- [258] Valaskova, V.; Baldrian, P. Degradation of cellulose and hemicelluloses by the brown rot fungus *Piptoporus betulinus*-production of extracellular enzymes and characterization of the major cellulases. *Microbiology*, **2006**, 152(12), 3613-22.
- [259] Osono, T.; Takeda, H. Fungal decomposition of *Abies* needle and *Betula* leaf litter. *Mycology*, **2006**, 98(2), 172-179.
- [260] Legonkova, O.A. Biotechnology of degradation of composite materials in different soils. *J. Balkan Tribological Association*, **2008**, 14(3), 405-412.
- [261] Kitamoto, H. Where are biodegradable plastic-degrading microorganisms? *BRAIN Techno News*, **2008**, 129, 1-6.

- [262] Sabev, H.; Handley, P.; Robson, G. Fungal colonization of soil-buried plasticized polyvinyl chloride (pPVC) and the impact of incorporated biocides. *Microbiology*, **2006**, 152(6), 1731-1739.
- [263] Nikaido, Y.; Oka, K.; Kondo, R. Biodegradation of soil vinyl alcohol polymers without dissolution or immersion in water. *Jpn. Kokai Tokkyo Koho*, **2007**, 12pp.
- [264] Nikaido, Y.; Kondo, R. Biodegradation poly(vinylalcohol)(PVA) film by Wood-rotting fungi. *Materiaru Raifu Gakkaishi*, **2007**, 19(1), 18-22.
- [265] Alariqi, S.A.S.; Kumar, A.P.; Rao, B.S.M.; Singh, R.P. Biodegradation of γ -sterilized biomedical polyolefins by synergistic mixtures of oligomeric stabilizers. *Polym. Degrad. Stabil.*, **2006**, 91(5), 1105-1116.
- [266] Coelho, N.S.; Almeida, Y.M. B.; Vinhas, G.M. The biodegradation of polyhydroxybutyrate-co-valerate/amphiprotic starch in the presence of microorganisms. *Polimeros: Ciencia e Tecnologia*, **2008**, 18(3), 270-276.
- [267] Bhatt, R.; Panchal, B.; Patel, K.; Sinha, V.K.; Trivedi, U. Synthesis, characterization, and biodegradation of carboxymethylchitosan-g-medium chain length polyhydroxyalkanoates. *J. Appl. Polym. Sci.*, **2008**, 110(2), 975-982.
- [268] Kamiya, M.; Asakawa, S.; Kimura, M. Molecular analysis of fungal communities of biodegradable plastics in two Japanese soils. *Soil Sci. Plant Nutrition*, **2007**, 53(5), 568-574.
- [269] Sakai, K.; Imai, T.; Yamanaka, H.; Moriyoshi, K.; Ohmoto, T.; Fujita, T.; Ohe, T. Distribution of biodegradable polymer-degrading microorganisms in the environment –in the case of poly(butylene succinate/adipate) and poly(lactic acid)s-. *Kagaku to Kogyo*, **2007**, 81(9), 453-457.
- [270] Sasek, V.; Vitasek, J.; Chromcova, D.; Prokopova, I.; Brozek, J.; Nahlik, J. Biodegradation of synthetic polymers by composting fungal treatment. *Folia Microbiol.*, **2006**, 51(5), 425-430.
- [271] Ishii, N.; Inoue, Y.; Shimada, K.; Tezuka, Y.; Mitomo, H.; Kasuya, K. Fungal degradation of Poly(ethylene succinate). *Polym. Degrad. Stabil.*, **2007**, 92(1), 44-52.
- [272] Ishii, N.; Inoue, Y.; Tagaya, T.; Mitomo, H.; Nagai, D.; Kasuya, K. Isolation and characterization of Poly(butylene succinate)-degrading fungi. *Polym. Degrad. Stabil.*, **2008**, 93(5), 883-888.
- [273] Hossain, Sk.M. Aerobic biodepolymerization studies of polyethylene using *Phanerochaete chrysosporium*. *Indian J. Environ. Protection*, **2006**, 26(11), 1006-1011.
- [274] Seneviratne, G.; Tennakoon, N.S.; Weerasekara, M.L.M.A.W.; Nandasena, K.A. Polyethylene biodegradation by a developed *Penicillium-Bacillus* biofilm. *Curr. Sci.*, **2006**, 90(1), 20-21.
- [275] Argolo, E.; Lins, M.C.M.; Palha, M.F.; Bastos de Almeida, Y.M.; Lima, M. Biodegradation of polymeric films by action of the *Talaromyces wortmanii* and *Phanerochaete chrysosporium* fungi. *Braz. Arch. Biol. Technol.*, **2006**, 49, 11-19.
- [276] Cosgrove, L.; McGeechan, P.L.; Robson, G.D.; Handley, P.S. Fungal communities associated with degradation of polyester polyurethane in soil. *Appl. Environ. Microbiol.*, **2007**, 73(18), 5817-24.
- [277] Friedrich, J.; Zalar, P.; Mohorcic, M.; Klun, U.; Krzan, A. Ability of fungi to degrade synthetic polymer nylon-6. *Chemosphere*, **2007**, 67(10), 2089-2095.
- [278] Labuzek, S.; Pajak, J.; Nowak, B.; Solga, M. Examination of toxicity of the products of polyester Bionolle degradation. *Polimery*, **2008**, 53(5), 384-389.
- [279] Trinh Tan, F.; Cooper, D.G.; Maric, M.; Nicell, J.A. Biodegradation of a synthetic *o*-polyester by aerobic mesophilic microorganisms. *Polym. Degrad. Stabil.*, **2008**, 93(8), 1479-1485.
- [280] Karthikeyan, C.; Sivakumar, N. Biodecolourization and biodegradation of anaerobically digested spent wash by an indigenous fungal culture. *Poll. Res.*, **2005**, 24(3), 567-570.
- [281] Begum, S.Y.A.; Noorjahan, C.M. Biodegradation of fertilizer industry effluent. *Asian J. Microbiol., Biotechnol. Environ. Sci.*, **2006**, 8(3), 585-588.
- [282] Kalyankar, N.V.; Kalyankar, S.N. Management of poultry feather waste using keratinolytic fungi. *Proc. Int. Conf. Solid Waste Technol. Manage.*, **2005**, 21, 1051-1058.
- [283] Beebi, Sk.K.; Lakshmi, M.V.V.C.; Sridevi, V.; Kusuma, M.P.; Rani, O.S. Biodegradation of ammonia using bacterial and fungal cultures. *Indian J. Environ. Protec.*, **2008**, 28(5), 430-433.
- [284] Yuan, H.L.; Yang, J.S.; Wang, F.Q.; Chen, W.X. Degradation and solubilisation of Chinese lignite by *Penicillium* sp. 6. *Appl. Biochem. Microbiol.*, **2006**, 42(1), 52-55.
- [285] Igbiginigie, E.E.; Aktins, S.; van Breugel, Y.; van Dyke, S.; Davies-Coleman, M.T.; Rose, P.D. Fungal biodegradation of hard coal by a newly reported isolate, *Neosartorya fischeri*. *Biotechnol. J.*, **2008**, 3(11), 1407-1416.
- [286] Hossain, Sk.M. Biological sugar industry wastewater treatment with fungus *Aspergillus niger*. *Indian J. Environ. Protec.*, **2008**, 28(6), 530-534.
- [287] Hossain, Sk.M.; Anantharaman, N. Aerobic biological pollution abatement of sugar industry wastewaters with *Phanerochaete chrysosporium*. *Process Plant Eng.*, **2007**, 24(4), 72-76.
- [288] Kalyankar, S.N.; Kalyankar, N.V. Biodegradation, management and utilization of fruit waste by aerobic composting. *Proc. Int. Conf. Solid Waste Technol. Manage.*, **2005**, 21, 1043-1050.
- [289] Huang, D.; Zeng, G.; Huang, G.; Jiang, X. Immobilized white-rot fungus preparation and its application in composting of garbage. *Faming Zhuanli Shenqing Gongkai Shuomingshu*, **2005**, 7 pp.
- [290] Subramanian, S.B.; Yan, S.; Tyagi, R.D.; Surampalli, R.Y. A new, pellet-forming fungal strain: its isolation, molecular identification, and performance for simultaneous sludge-solids reduction, flocculation, and dewatering. *Water Environ. Res. : a research publication of the Water Environment Federation*, **2008**, 80(9), 840-52.
- [291] Rema, T.; Srinivasan, S.V.; Chitra, K.; Umamaheswari, B.; Balakameswari, K.S.; Ravindranath, E.; Suthanthararajan, R.; Rajamani, S. Removal of color from secondary treated tannery effluent using *Trametes versicolor*. *Indian J. Environ. Protec.*, **2005**, 25(9), 784-787.
- [292] Strobel, G.; Dirkse, E.; Ezra, D.; Castillo, U.; Phillips, B. A method of using endophytic fungi to decontaminate and decompose human and animal wastes. *PCT Int. Appl.*, **2005**, p. 40.
- [293] Chen, S.; Zhang, X.; Xu, H.; Xie, J. Biodegradation of DDNP wastewater pretreated by micro-electrolysis using white rot fungus. *Huanjing Wuran Yu Fangzhi*, **2006**, 28(2), 143-146.
- [294] Bhatt, M.; Zhao, J.; Halasz, A.; Hawari, J. Biodegradation of hexahydro-1,3,5-trinitro-1,3,5-triazine by novel fungi isolated from unexploded ordnance contaminated marine sediment. *J. Ind. Microbiol. Biotechnol.*, **2006**, 33(10), 850-858.
- [295] Lin, J.; Tang, W.; Lin, J.; Zhou, S.; Zheng, P. Degradation of royal demolition explosive by white-rot-fungi. *Zhongguo Jishui Paishui*, **2006**, 22(7), 74-77.
- [296] Gao, X.; Zhang, X.; Zhou, S. Biodegradation of RDX wastewater by white rot fungi. *Jiangsu Huagong*, **2005**, 33(6), 56-58.
- [297] Fournier, D.; Monteil-Rivera, F.; Halasz, A.; Bhatt, M.; Hawari, J. Degradation of CL-20 by white-rot fungi. *Chemosphere*, **2006**, 63(1), 175-181.